

Review

Open Access

Using non-invasive brain stimulation to augment motor training-induced plasticity

Nadia Bolognini^{1,2,3}, Alvaro Pascual-Leone^{1,3} and Felipe Fregni*¹

Address: ¹Berenson-Allen Center for Noninvasive Brain Stimulation, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, USA, ²Department of Psychology, University of Milano Bicocca, Milano, Italy and ³Institut Guttmann de Neurorehabilitació, Universitat Autònoma de Barcelona, Barcelona, Spain

Email: Nadia Bolognini - nadia.bolognini@unimib.it; Alvaro Pascual-Leone - apleone@bidmc.harvard.edu; Felipe Fregni* - ffregni@bidmc.harvard.edu

* Corresponding author

Published: 17 March 2009

Received: 17 November 2008

Accepted: 17 March 2009

Journal of NeuroEngineering and Rehabilitation 2009, **6**:8 doi:10.1186/1743-0003-6-8

This article is available from: <http://www.jneuroengrehab.com/content/6/1/8>

© 2009 Bolognini et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Therapies for motor recovery after stroke or traumatic brain injury are still not satisfactory. To date the best approach seems to be the intensive physical therapy. However the results are limited and functional gains are often minimal. The goal of motor training is to minimize functional disability and optimize functional motor recovery. This is thought to be achieved by modulation of plastic changes in the brain. Therefore, adjunct interventions that can augment the response of the motor system to the behavioural training might be useful to enhance the therapy-induced recovery in neurological populations. In this context, noninvasive brain stimulation appears to be an interesting option as an add-on intervention to standard physical therapies. Two non-invasive methods of inducing electrical currents into the brain have proved to be promising for inducing long-lasting plastic changes in motor systems: transcranial magnetic stimulation (TMS) and transcranial direct current stimulation (tDCS). These techniques represent powerful methods for priming cortical excitability for a subsequent motor task, demand, or stimulation. Thus, their mutual use can optimize the plastic changes induced by motor practice, leading to more remarkable and outlasting clinical gains in rehabilitation. In this review we discuss how these techniques can enhance the effects of a behavioural intervention and the clinical evidence to date.

Introduction

Motor impairments following stroke or traumatic brain injury (TBI) are the leading cause of disability in adults. More than 69% of all stroke survivors experience lasting functional motor impairments in the upper limbs and approximately 56% continue to complain of marked hemiparesis as long as 5 years post-stroke [1-5]. Such losses in function can severely impact quality of life and the functional independence in numerous activities of daily living [4,5]. Similarly, after TBI, fine and gross motor deficits are frequently observed. Complementary impair-

ments such as ataxia, movement disorders and vestibular impairments, can also potentially affect motor functioning in TBI. Moreover, other factors such as multiple trauma, resulting in musculo-skeletal and peripheral nervous system injury, also complicate the recovery of motor functions in these patients [6].

Although some degree of recovery may occur spontaneously, there is strong evidence that intensive practice is essential in order to substantially promote motor recovery [7-9]. As shown by several neurobehavioral discoveries in

animals and humans, such experience-dependent change can occur at multiple levels of the central nervous system, from the molecular, to the synaptic level of cortical maps and large-scale neural networks [10,11].

Standard motor therapies involve different approaches aimed at improving motor functions by minimising impairment or developing suitable adaptation strategies. For instance, neurofacilitation techniques are aimed at retraining motor control by promoting normal (recruitment of paretic muscles) while discouraging abnormal movement or muscle tone. Different facilitation approaches have been developed, including cutaneous/proprioceptive, weight bearing, proximal pre-innervation, and contralateral pre-innervation [12]. Task-specific training is aimed at improving skill in performing selected movement or functional tasks: examples of this type of treatment are index finger tracking [13] or the combination of task-specific motor training with the inhibition of ipsilesional sensorimotor cortex representation of the paretic upper arm by local anaesthesia [14]. Finally, task-oriented training aimed at retraining functional tasks by taking into account the interplay of different systems is another possible approach. For example, constrained-induced movement therapy (CIMT) combines intensive physical practice using the affected upper limb with restricted use of the unaffected upper limb in order to prevent its habitual compensatory utilization [15]. Bilateral arm training is instead based on the phenomenon of interlimb coupling, in which the movement patterns of the arms are similar when moving simultaneously [16,17]. Ongoing studies indicate that even mere action observation, activating the same cortical motor areas that are involved in the performance of the observed actions (i.e. action observation/execution matching system) can lead to a reorganization of the motor system resulting in an improvement of motor functions [18,19]. Other treatments have focused on the use of robotics [20,21], EMG-triggered stimulation [22], and motor imagery [23] (see for a review [24]).

Although there is little doubt that behavioural motor therapy clearly plays a role in promoting contra- and ipsilesional plastic changes after stroke, the functional outcomes are often of limited practical significance and after completing standard rehabilitation approximately 50–60% of patients still exhibit some degree of motor impairment and require at least partial assistance in activities of day living [24,25]. Similarly, the efficacy of the majority of standard motor interventions for promoting recovery after TBI is supported by rather limited evidence [6]. Therefore, investigation of other approaches to promote the recovery of motor impairments is essential. In this context, noninvasive brain stimulation (NIBS) appears to be an interesting option [26]. Transcranial Magnetic Stim-

ulation (TMS) is delivered to the brain by passing a strong brief electrical current through an insulated wire coil placed on the skull. Current generates a transient magnetic field, which in turn, if the coil is held over the subjects head, induces a secondary current in the brain that is capable of depolarising neurons. Depending on the frequency, duration of the stimulation, the shape of the coil and the strength of the magnetic field, TMS can activate or suppress activity in cortical regions [27]. Another method of non-invasive brain stimulation is transcranial Direct Current Stimulation (tDCS) which delivers weak polarizing direct currents to the cortex via two electrodes placed on the scalp: an active electrode is placed on the site overlying the cortical target, and a reference electrode is usually placed over the contralateral supraorbital area or in a non-cephalic region. tDCS acts by inducing sustained changes in neural cell membrane potential: cathodal tDCS leads to brain hyperpolarization (inhibition), whereas anodal results in brain depolarization (excitation) [28,29]. Differences between tDCS and TMS include presumed mechanisms of action, with TMS acting as neuro-stimulator and tDCS as neuro-modulator. Moreover, TMS has better spatial and temporal resolution, TMS protocols are better established, but tDCS has the advantage to be easier to use in double-blind or sham-controlled studies [30] and easier to apply concurrently with behavioural tasks (for discussion of these methods, similarities and differences, see the review by Wagner et al. [31]). Despite their differences, both TMS and tDCS can induce long-term after-effects on cortical excitability that may translate into behavioural impacts that can last for months [32-35]. These long-term after-effects are believed to engage mechanisms of neural plasticity, rendering these techniques ideally suited to promote motor recovery particularly when combined with suitable behavioural interventions (for review, see [26,36,37]).

To date, two approaches have been tested. They are based on a model of interhemispheric rivalry between motor areas in the damaged and undamaged (intact) hemispheres. In essence, the model proposes that motor deficits are due to reduced output from the damaged hemisphere and excess inhibition of the damaged hemisphere from the intact hemisphere [26,38]. Thus, improvement may be possible by either up-regulating excitability of the lesioned motor cortex or down-regulating excitability in the intact motor cortex [26]. Enhancement of excitability can be achieved with either high frequency rTMS and anodal-tDCS. Suppression of excitability can be accomplished with either low-frequency rTMS and cathodal-tDCS. A growing body of evidence from small clinical trials has demonstrated the efficacy of both approaches to induce considerable changes on cortical excitability, which often correlate with relevant clinical gains in motor functions. However, most studies to date

have examined the effects of NIBS without coupling it with any specific behavioural, physical or occupational therapy, and the functional benefits are often limited, inducing about 10–20% functional improvement in some single-session and longer-term therapeutic trials [37]. This is probably a suboptimal approach, as NIBS activates neural circuits in a non-specific way. Therefore given that NIBS and motor training are thought to share synergistic impacts on synaptic and network plasticity an emerging field of research is focusing on the possibility of coupling both therapies in order to achieve additive practical impact. The underlying principle of this approach is that practice of a motor task may be more effective at using the (surviving) neural mechanisms sub-serving training-dependent plastic changes if pertinent areas of the cortex are facilitated [38]. In addition, motor training can guide the activation of specific neural networks associated with the desired behavior. Considering for instance that many of the spontaneous plastic changes induced by a stroke, including phenomena of hyperexcitability, diminish after a few months [39-41], the therapeutic window of potential plastic changes for motor recovery seems to be limited. NIBS might be helpful to prolong this therapeutic window thus offering a greater opportunity for suitable physical and occupational therapies to promote functional recovery. Although preliminary, there is some recent encouraging evidence supporting the clinical validity of this approach.

Mechanisms of NIBS to induce neuroplasticity

After a stroke affecting the motor cortex, cortical excitability is generally decreased in the affected primary motor cortex relative to the unaffected motor cortex. This might result from a shift in interhemispheric interactions, with increased transcallosal inhibition from the intact to the damaged motor cortex [41,42]. In this scenario, TMS and tDCS applied over the intact hemisphere allow safe cortical stimulation in humans in order to promote restoration of activity across bihemispheric neural networks and guidance towards more-adaptive plasticity [26,43].

TMS uses a rapidly changing magnetic field to induce electric currents via electromagnetic induction. A very brief high-intensity electric current is passed through a wire coil held over the scalp, this generates a magnetic field pulse which passes relatively unimpeded through the layers of tissue and bone and reaches the brain where secondary currents are induced. These secondary currents are induced in a plane parallel to the plane of the stimulation coil, which typically is held tangentially to the scalp, over the subject's head. Current direction and electric field distribution depend on output pulse shape of the stimulator and coil geometry respectively. The secondary current can be sufficient to depolarize cortical neurons, directly at their axon hillock or indirectly via depolarization of

interneurons. Exactly which neural elements are activated by TMS and the mechanisms of neuronal stimulation remains unclear and might be variable across different brain areas and different subjects [27]. We know that when TMS is delivered over the primary motor cortex with adequate intensity, it induces efferent volleys along the corticospinal pathway [44]. Crucially, the therapeutic relevance of this technique is due to the long-term effects that occur after repeated stimulation. TMS delivered in a repetitive mode (rTMS) can indeed modulate cortical excitability beyond the duration of the rTMS trains themselves [45]. Depending on rTMS parameters, long lasting suppression or facilitation of cortical excitability can be induced: low-frequency rTMS (≤ 1 Hz) usually results in decreased cortical excitability [46], whereas at higher frequencies (>1 Hz) cortical excitability is usually increased [45]. It should however be noted, that this is an average effect across individuals, and yet there is substantial inter-individual variability as well as intra-individual variability depending on the timing and exact location of stimulation [47,48].

In promoting stroke recovery, both, high frequency rTMS and low frequency rTMS have been tested and appear promising. For instance, Takeuchi et al. [49] and Fregni et al. [32] applied low-frequency rTMS to suppress activity in the contralesional (undamaged) hemisphere in chronic stroke patients: this suppressive protocol proved to be effective in reducing the transcallosal inhibition from the intact to the affected motor cortex [49] and increasing excitability of the lesioned motor cortex [32]. On the other hand, up-regulating the excitability of the lesioned M1 can also be successful. Talelli et al. (2007) reported that a single session of excitatory intermitted theta burst stimulation (TBS), consisting in delivering 3 pulses at 50 Hz, repeated at a rate of 5 Hz, increased MEP amplitude on the stroke side, with additional transiently improvement of motor behaviour [50]. By contrast, in the same study, continuous TBS of the unaffected motor cortex, which like low frequency rTMS suppresses excitability, did not change motor behaviour or the electrophysiology of the paretic hands [50]. Di Lazzaro et al (2008) obtained slightly different results. They showed that in acute stroke patients both intermittent TBS over the stroke hemisphere and continuous TBS over the intact hemisphere enhanced the excitability of the lesioned motor cortex and resulted in a functional benefit [51].

Despite these promising results, some limitations of TMS need to be noted. Critically, after stroke, there is a change in the local anatomy and the lesion evolves in time to formation of scar tissue and, particularly in the case of cortical damage, larger cerebrospinal fluid spaces. Because the conductance of cerebrospinal fluid (CSF) is 4 to 10 times higher than that of brain tissue, scar formation and larger

CSF spaces modify the geometry and magnitude of the electric field induced by rTMS, and stimulation of the lesioned hemisphere can become difficult to predict unless careful modelling is done [52].

The mechanisms underlying long-term effects of TMS are incompletely understood, but they could be analogous to long-term potentiation (LTP) or depression (LTD) seen in the hippocampus after repeated activation of synaptic pathways [53-55]). In addition, modulation of neurotransmitter levels seems to be a contributing factor. The neurotransmitter systems involved include the inhibitory GABAergic system [56-58] as well as the excitatory glutamatergic system with activation of NMDA receptors [57]. TMS may result in changes in endogenous neurotransmitters (GABA and glutamate) and neuromodulators (DA, NE, 5-HT, ACh) which play a pivotal role in the regulation of the neuronal activity in the cerebral cortex (for review, [59]). A focal increase of dopamine in the striatum was indeed demonstrated in healthy human after sub-threshold 10 Hz rTMS applied to the ipsilateral primary motor cortex [60] or dorsolateral prefrontal cortex [61].

Another candidate mechanism by which rTMS may exert persistent effects is through gene induction. Actually, rTMS can modulate the expression of immediate early genes [62-64]. A single rTMS train increased *c-fos* mRNA in the paraventricular nucleus of the thalamus and, although to a lesser extent, in the frontal and cingulate cortices [64]. A longer treatment protocol (up to 14 daily sessions) could even induce an increase in *c-fos* mRNA in the parietal cortex of rodents [63] and an enhancement of BDNF mRNA in the hippocampus, the parietal and piriform cortices [65]. As suggested, BDNF is a neurotrophic factor that is critically linked to the neuroplastic changes [66] and might serve to index neuroplastic effects induced by rTMS [67].

The other main method of NIBS, tDCS, is a form of brain polarization that uses prolonged low-intensity electric current (1–2 mA) delivered to two large electrodes (usually 5 × 7 cm or 5 × 5 cm) to the scalp. To stimulate the primary motor cortex, usually one electrode is placed on the scalp over M1 and the other on the contralateral supraorbital area [68]. Alternatively, the reference electrode can be placed on the shoulder or another extra-cranial location. Reminiscent of the effects of repetitive TMS, tDCS can up- or down-regulate neural activity in the stimulated regions. Increased excitability of the underlying neurons occurs with anodal stimulation, while decreased excitability is seen after cathodal stimulation. With only 13 minutes of tDCS stimulation, effects on neural excitability outlasts the period of stimulation by up to 90 minutes [69]. In fact, the after-effects of tDCS appear greater than those induced by synchronous rTMS [68,70]. How-

ever, TBS or other, more sophisticated, asynchronous rTMS trains may significantly enhance and prolong the modulatory effects.

Again reminiscent of the effects of rTMS, tDCS-induced changes in cortical excitability are associated with changes in the excitability of inhibitory and facilitatory intracortical circuits: whereas anodal tDCS results in decreased intracortical inhibition and increased intracortical facilitation, cathodal stimulation induces opposite effects. In patients with chronic strokes, either anodal tDCS delivered to the lesioned M1 or cathodal tDCS delivered to the contralesional hemisphere can result in an improvement in motor functions [71-73].

tDCS does not stimulate axons and cause them to discharge action potentials, as TMS does. Rather, it most likely targets neuronal signalling by manipulating ion channels or by shifting electrical gradients which influence the electrical balance of ions inside and outside of the neural membrane; thus modulating the resting membrane threshold. Apart from membrane potential changes, chemical neurotransmission, either pre- or post-synaptically, may play a role in tDCS effects [74]. Some studies have aimed to clarify the cellular mechanisms of tDCS over the motor cortex [29,74]. For instance, the effects of the sodium channel blocker carbamazepine, the calcium channel blocker flunarizine and the NMDA-receptor antagonist dextromethorphan on tDCS-elicited motor cortex excitability changes were tested in healthy human subjects. Carbamazepine selectively eliminated the excitability enhancement induced by anodal stimulation during and after tDCS. Flunarizine resulted in similar changes. Antagonizing NMDA receptors did not alter current-generated excitability changes during stimulation, but prevented the formation of after-effects independent of their direction. Therefore, authors concluded that cortical excitability shifts induced during tDCS in humans appear to depend on membrane polarization, thus, modulating the conductance of sodium and calcium channels. In addition, the after-effects seem to be NMDA-receptor dependent. Recently, it was demonstrated that d-cycloserine, a partial NMDA-agonist, selectively potentiates the duration of motor cortical excitability enhancements induced by anodal tDCS [75]. Additionally, it was also suggested that the after-effects of cathodal tDCS include nonsynaptic mechanisms based on changes in neuronal membrane function [76]. Long term effects induced by tDCS may include built-up of new synapses, with mechanism of LTP and LTD critically involved. The glutamatergic system, in particular NMDA receptors [77], seems to be necessary for induction and maintenance of neuroplastic after-effect excitability enhancement and reduction induced by tDCS [74].

Mechanisms of action of motor training in inducing plastic changes

After brain damage, there is substantial recovery with clearly delineated dynamics, resulting in a faster recovery in the acute and subacute stages, gradually levelling off as time progresses. In addition activity-dependent long-term modification of synaptic efficacy is associated with information storage in neural networks [78]. In fact, Neural plasticity changes evolve from the Hebbian synapse rule that states that individual synaptic junctions respond to activity (use) and inactivity (disuse) [79].

Motor training can promote plastic changes in injured motor networks even in a chronic stage of illness. However, simple interventions such as repetitive movement practice fail to induce profound plastic changes [80]. It appears that skill learning must be present to promote cortical plasticity [81]. In fact, most of the recovery of function after a stroke may represent actual relearning of the skills with the injured brain. Recovery mediated by training, like learning in healthy subjects, is usually task-specific and it differs from processes involved in compensation: whereas recovery of motor functions requires the recruitment of brain areas to generate commands to the same muscles as were used before the injury, compensation is instead based on the use of alternative muscles to accomplish the task goal [82]. Motor learning will lead first to strengthening of existing neural pathways, and second, to new functional or structural changes and thus expression of neuroplasticity [8].

The main mechanism underlying this relearning process after stroke involves shifts of distributed contributions across a specific neural network. Investigations in adult animals have revealed that motor learning can promote a plastic reorganization of motor maps in M1 with the representations of specific movements used to perform the motor task selectively expanding in the motor cortex at the expense of other areas not used for forelimb representations [10]. Similar results have been obtained in humans. For instance, the acquisition of new fine motor program induces an enlargement of the cortical motor areas targeting the muscles involved in the task, with an additional decrement of the activation threshold, as measured by means of TMS. Such map expansions parallel improvements in motor performance [83]. These results indicate that the cortex has the potential for rapid and large-scale functional changes in response to motor skill learning. One important issue is that an enlargement of a given neural network occurs at the cost of modifying another network and therefore with the theoretical risk of decreasing performance in another task. To date, this theoretical concern does not seem to cause any significant impairments in stroke subjects receiving intensive motor training.

Evidence for a long-term alteration in brain function associated with a therapy-induced motor recovery in neurological populations has also been provided. For instance, constraint-induced movement therapy (CIT) can significantly change cortical excitability measured by TMS in both affected and unaffected hemispheres. More specifically, CIT can result in an enlargement of the motor output map in the affected hemisphere, which is associated with a greatly improved motor performance of the paretic limb. A shift on the center of gravity of the output map in the affected hemisphere was also observed, indicating the recruitment of adjacent brain areas. Follow-up examinations up to 6 months after treatment showed that motor performance remained at a high level, whereas the cortical area sizes in the two hemispheres became almost identical, representing a return of the balance of excitability between the two hemispheres toward a normal condition [84]. These results are in line with PET and fMRI studies in recovered stroke patients showing that plastic changes taking place within the ipsilateral, noninfarcted hemisphere might contribute to the restitution of motor function [7,85-87]. A recent meta-analysis further underlines the positive impact of motor rehabilitation for the upper extremities, showing that practice-dependent recruitment of the ipsilesional hemisphere induces clear functional motor gains [88]. Increased engagement of the damaged hemisphere is expressed by either an increase in the area of the brain subserving the paretic arm movement, as shown by brain imaging techniques, and by greater signal strengths of physiological-functional measures (MEPs) within the sensorimotor cortex of the lesioned hemisphere [88].

Although motor training can lead to neurofunctional adaptation within a matter of minutes [89], long-term representational changes may take days [83] or weeks of practice [90]. Rapid changes are bound to be reflected in a less specific remodelling of network activity [91]. Instead, enduring change is reflected in, for example, augmented dendritic branching [92] and synaptogenesis [93], possibly provoked by specific gene induction [94,95]. Ultimately these processes result in an increase in the efficacy of synaptic transmission [96]. In strict analogy with the NIBS-induced after-effects, NMDA receptor activation and GABAergic inhibition are likely mechanisms operating in use-dependent plasticity in the intact human motor cortex and point to similarities in the mechanisms underlying this form of plasticity and long-term potentiation (LTP) [97]. LTP is associated with the proliferation of dendritic spines [98]. This morphologic change has been even found in homologous cortex opposite from the site of an experimental sensorimotor cortical lesion when the unaffected limb works to compensate for the paretic one [99]. This evidence suggests that the synaptic strength of horizontal connections in the motor cortex are modifiable

and may provide a substrate for altering the topography of cortical motor maps during physical intervention based on motor learning.

Combination of NIBS with motor training to enhance neuroplasticity and behavioural changes

As we have seen, motor learning and NIBS may share similar mechanisms of action for inducing neuroplastic changes in the human cortex. One possible conjecture then, is that their combination might mutually maximise their individual effects. Since learning processes are accompanied by cortical excitability shifts and by changes of synaptic efficacy and considering that the after-effect of NIBS is NMDA-receptor dependent, there is a possibility that cortical excitability changes induced by cortical stimulation could interact with ongoing motor learning process, improving learning-related NMDA-receptor strengthening. It is noteworthy that synaptic plasticity is bidirectional [100]. The basic idea is that the ongoing state of the cortex at the time of physical therapy can reinforce the long-term effects induced by motor practice. If so, the rationale of coupling NIBS and motor therapy is that it is possible to enhance or depress the response of a neural network to a form of stimulation, e.g. motor training, by previous priming it with a different form of stimulation, e.g. NIBS (and vice versa). Some experimental studies provide preliminary support to this hypothesis.

Animal studies using direct repetitive electric stimulation (ES) of the cortex – a technique that mimics rTMS and can alter cortical excitability as measured by cortical spreading depression (CSD) [101] have supported the importance of priming brain activity. CSD is an indicator of cortical excitability [102] characterised by alterations in cerebrocortical ion homeostasis in response to the direct stimulation of brain tissue. The alterations result in a wave of neuronal excitation propagating through the cortex followed by transient inhibition. It has been found that when active or sham 1 Hz ES was applied to Wistar rats preconditioned with active, sham or cathodal tDCS, a pattern suggestive of homeostatic mechanisms emerged [101]: 1 Hz ES that was applied alone or was preceded by cathodal tDCS, reduced CSD velocity whereas anodal tDCS followed by 1 Hz ES increased CSD velocity. Homeostatic effects have also been found in the effects of tDCS on paired associative stimulation (PAS) of human motor cortex [103] or by preconditioning of rTMS with tDCS [104]. However there is a fundamental difference when coupling two techniques of neuromodulation vs. coupling neuromodulation techniques with motor training; the latter might be better as it can focalize the effects to specific networks. In fact, several studies have explored the influence of coupling learning tasks with NIBS on motor and cognitive functions in healthy subjects. In one example, TMS synchronously applied to a motor cortex

engaged in a motor learning task was shown to be effective in enhancing use-dependent plasticity. Healthy volunteers were studied in different sessions: training alone, training with synchronous application of TMS to the motor cortex contralateral or ipsilateral to the training hand, and training with asynchronous TMS. It was found that the longevity of use-dependent plasticity was significantly enhanced only by TMS applied in synchrony to the cortex contralateral to the training hand [105]. Carey et al. (2006) have obtained, however, different results: investigation of the effects of motor learning training, consisting in finger tracking with the right hand, unexpectedly showed that 1 Hz rTMS interfered transiently with motor performance when applied ipsilateral to the training hand but it had no effect when applied contralaterally [106]. In another tDCS study [107], the excitability of MT+/V5 and M1 was increased or decreased by anodal or cathodal tDCS while subjects were learning a visually guided manual tracking task. Accuracy of tracking movements was increased significantly by anodal stimulation, whereas cathodal stimulation had no significant effect on visual learning. Interestingly, the positive effect of anodal tDCS was restricted to the learning phase, suggesting a highly specific effect of the stimulation. Similar results were demonstrated for implicit motor learning in healthy human subjects [108], and in addition a recent study showed the beneficial effects of anodal tDCS over the posterior part of the left peri-sylvian area on language learning [109]. One important conclusion is that the effects are dependent on the site of stimulation, task and parameters of stimulation, therefore making difficult to generalize conclusions on these studies and also opening the possibility to induce detrimental effects when coupling these two intervention methods; therefore, it is critical to study the combination of these techniques before using it in clinical practice.

Originally, encouraging results have been found in animal investigations. Seminal experiments in animals have shown that coupled forced use of the paretic hand with implanted electrical stimulation to the ipsilesional M1 lead to significant behavioural improvements with large-scale expansions of the hand representation into areas previously representing proximal forelimb movements [80,110]. In a similar way, a recent prospective, randomized, multicenter study showed that in chronic stroke intensive motor therapy combined with invasive epidural electrode is associated with a significant improvement in motor function [111].

Although investigation with NIBS is still at the beginning, there are some very promising preliminary results. Khedr et al. (2005) have explored the effects of rTMS in patients with acute ischemic stroke as an add-on intervention to standard physical and drug therapies. rTMS was applied

over the M1 of the stroke hemisphere for 10 days. rTMS consisted in ten 10-second trains of 3-Hz stimulation with 50 seconds between each train [112]. The motor treatment consisted in the passive limb manipulation, increasing by the end of the first week to more active movements if patients improved function. Treatment effects were measured with clinical scales and neurophysiological measurements – i.e., resting motor threshold (RMT) of healthy side, and motor evoked potentials (MEPs) of healthy and hemiplegic sides. On every scale, patients' motor scores in the active rTMS group had a significant greater improvement as compared with sham rTMS, leading to a higher percentage of independent patients and a higher percentage of patients having only mild disability by the time of the follow-up assessment, after 10 days from the end of the treatment. However, no effect was seen in patients with massive middle cerebral artery infarcts. 14 out of 21 patients in the real rTMS group recovered MEPs; although MEPs tended to improve more in the real rTMS group, this was not significantly different from the sham group. In addition, no correlation between clinical recovery and changes in MEP was found [112].

In other clinical trial [113], patients with chronic hemiparetic stroke practiced a complex, sequential finger motor task using their paretic fingers either after receiving high-frequency (10 Hz, repeated 8 times) or sham rTMS over the primary motor cortex (M1) of the damaged hemisphere. Changes in the behavior and corticomotor excitability before and after the intervention were examined by measuring the movement accuracy, the movement time, and the MEP amplitude. The authors found that rTMS induced a significantly larger increase in the MEP amplitude than sham rTMS; this corticomotor excitability change was associated with enhanced motor skill acquisition.

Rather than trying to enhance the cortical excitability of the damaged motor cortex, Takeuchi et al (2008) explored the effect of inhibiting the contralesional motor cortex in chronic patients [114]. Patients were randomly assigned to receive either a sub-threshold rTMS over the unaffected hemisphere (1 Hz, 25 minutes) or sham stimulation and all patients performed a pinching task after stimulation. Compared with sham stimulation, rTMS induced an increase in the excitability of the injured motor cortex and an improvement in acceleration of the affected hand. The effect of motor training on pinch force was also enhanced by rTMS. Such improvement was stable at the follow-up examination, one week after the intervention [114]. Another study [115] assessed the efficacy of low-frequency 1 Hz rTMS combined with voluntary muscle contraction (VMC) on corticospinal transmission, muscle function, and purposeful movement early after stroke. rTMS consisted of 5 blocks of 200 1-Hz stimuli (using an

interblock interval of 3 minutes), applied to the lesioned hemisphere. The treatment was given for 8 working days. The motor training task in this study was VMC – the paretic elbow was repeatedly flexed/extended for 5 minutes. The main finding was that in patients who underwent the rTMS combined with VMC, motor-evoked potential frequency increased 14% for biceps and 20% for triceps; whereas, with Placebo rTMS plus Placebo VMC, motor-evoked potential frequency decreased 12% for biceps and 6% for triceps.

Negative findings have also been reported. A recent study indeed did not prove the usefulness of combining rTMS stimulation with a standard motor therapy [116]. Here, chronic stroke patients undergoing ten days of constraint-induced therapy (CIT) for upper-limb hemiparesis, which was combined with 20 Hz rTMS (stimulus train duration of 2 secs, intertrain interval of 28 secs.) or with sham rTMS of the affected M1. Primary outcome measures to assess change in upper-extremity function were the Wolf Motor Function Test (WMFT) [117] and the Motor Activity Log (MAL)-Amount [118]. Secondary outcome measures included the MAL-How Well and the Box and Block Test (BBT) [119] and MEP threshold. The results showed that, regardless of the rTMS intervention, participants demonstrated significant gains on the primary outcome measures and on secondary outcome measures, further supporting the efficacy of CIT. Indeed, although a significant decrease in motor threshold for subjects receiving rTMS was found, which was not observed after sham rTMS, this increase in the excitability of the motor system did not translate into a clinically evident outcome. Ceiling effects and outcome measures might have contributed to these findings.

tDCS is another technique associated with a significant beneficial effect on motor recovery after stroke. Although its beneficial effects on motor function have been shown by several small studies [26,32,43,71,72,120], actually there are very few clinical trials of the potential adjuvant of this technique to physical therapy. Yet, tDCS might be a more suitable tool to enhance the effects of motor training as it offers several advantages as compared with TMS in a rehabilitation setting. For instance, whereas rTMS has to be delivered in a off-line paradigm and it usually precedes the behavioural intervention, the portable use of tDCS allow to deliver the cortical stimulation during the motor training. Moreover, tDCS modulatory effects last longer as compared to rTMS – for example, 13 minutes of stimulation changes brain excitability for up to 90 minutes [69]. Finally, due to its physiological effect on the membrane resting potential, tDCS could to be more appropriate for priming motor neural network for subsequent stimulation with tDCS. Hummel et al (2005) have recently explored the effects of tDCS on skilled motor functions in chronic stroke patients [121]. Anodal tDCS

was delivered for 20 min to the affected hemisphere during the execution of the Jebsen-Taylor Hand Function Test (JTT), a widely used assessment of functional hand motor skills. Active anodal tDCS was associated with improvements in motor function of the paretic hand. The magnitude of tDCS-induced improvement in JTT was approximately 11.75% (+/- 3.61%) and persisted for more than 25 min after the stimulation ended. However, patients' performance returned to baseline levels after 10 days of the end of stimulation [72]. In another preliminary report in chronic stroke, anodal tDCS (1.5 mA) was combined with robot-assisted arm training (AT) [122]. Over six weeks, patients received 30 sessions of 7 min tDCS integrated into 20 min of AT. Arm function of three out of ten patients (two of them with a subcortical lesion) improved significantly, as measured by the Fugl-Meyer motor score. In the remaining seven patients, all with cortical lesions, arm function changed little. However, this study lacked an adequate control group and it included a small number of patients, who were still in the phase of spontaneous recovery; therefore no definite conclusions can be made.

Overall, the data discussed above provide some encouraging information supporting the proposal that NIBS might optimize the effect of standard physical therapy under certain circumstances. Beyond the obvious need for further clinical trials to corroborate the validity of this approach, attention must be directed in understanding the optimal way to combine motor training with NIBS. Crucially the next step is to determine the best parameters required to optimize the conditioning effects of NIBS on motor therapy, as well as the exact temporal window during which NIBS can be delivered in order to modulate brain plasticity and enhance the effects of the motor training.

How brain stimulation should be used in combination with motor training – methods of optimizing functional improvements

Given the limited number of clinical trials that have assessed the efficacy of combining NIBS with physical therapy, any prediction of the clinical utility of this approach remains speculative. Although further investigations are needed to make any relevant clinical consideration, some reflections can be delineated in order to make an optimal use of this approach in the near future. So far, the best option in order to optimize the effects of coupling NIBS and motor therapy still needs to be explored but it likely may depend on different factors, as the stages of illness (e.g. acute versus chronic), the type of motor training, the site of stimulation, the timing of stimulation in relation to physical intervention, baseline cortical activity and the technique of NIBS used. An essential issue to take into account when applying these NIBS protocols to a damaged human brain is related to the concept of homeostasis

– that is the human's brain ability to regulate changes in synaptic plasticity as to avoid drastic changes in its function. Therefore homeostasis is likely to respond definitively and forcefully to artificial and functionally non-specific changes in network activity such as those probably induced by NIBS [123]. Homeostatic plasticity (i.e., the dependency of the amount and direction of the obtainable plasticity from the baseline of a neuronal network) is increasingly recognized as regulatory mechanism for keeping neuronal modifications within a reasonable physiological range. Homeostasis provides a means for neurons and circuits to maintain stable functions in the face of perturbations such as activity-dependent changes in synapse number or strength [124]. In this regard, recent experimental works emphasize the importance of homeostatic plasticity as a means to prevent destabilization of neuronal networks that could operate in neurorehabilitative settings [124,125]. In particular, as advised by Thickbroom (2007), the influence of homeostatic mechanisms cannot be overlooked either during or after NIBS interventions: homeostatic mechanisms could be a crucial factor in repeat interventions, as are sometimes employed in NIBS protocols, or for intervention protocols of longer duration in which they may begin to act during the intervention itself. They could be one of the main factors that limit the magnitude and duration of post-TMS effects. For instance, NIBS could evoke compensatory regulatory mechanisms, which are a part of the process of maintaining normal brain function. On the other hand, activity-dependent forms of plasticity, even those incorporating LTP and LTD mechanisms, are inherently unstable due to positive feedback [123]. Thus, the successful implementation of NIBS as adjuvant strategy to physical therapy should rely on an improved understanding of the underlying plastic mechanisms and their functional interaction with activity-induced plasticity. For instance, a challenging issue is the time of the NIBS intervention relative to the motor task. As seen above, when combining to a motor training, so far NIBS has been usually delivered just before the task. However, functional therapies could in principle be implemented at different phases in conjunction with a NIBS intervention. NIBS preceding the motor training could potentially prime functional networks for the physical intervention. Instead, NIBS simultaneously applied during a behavioural intervention might preferentially interact with the networks selectively recruited by the ongoing task. Even the application of NIBS after motor training could be a potential choice; the underlying rationale of this approach is that, after the modulation induced by the motor therapy, a further modulation of cortical excitability might selectively build up the activity-dependent activation of a given network and promote its functional stabilization. It is not completely unlikely that even after excitability has returned to baseline, perhaps due to homeostatic regulation, NIBS could still be func-

tionally beneficial [123]. Here, NIBS could be influential for driving longer-term consolidation of new network patterns. The choice of the more suitable time window for NIBS intervention likely needs careful examination in order to exclude maladaptive cortical responses, which could interfere with or even suppress the effects of the behavioural therapy. For instance, an excitability modulation induced by tDCS during the performance of a motor task might be best suited to improve motor learning than tDCS administered prior of learning or motor behaviour [126,127]. This is because, during tDCS not only NMDA receptors, but also calcium channels are modulated, while the after-effects of tDCS are achieved by modifications of NMDA receptors alone [29]. Since intracellular calcium concentration is important for LTP induction [128] enhanced transmembrane calcium conduction, as probably achieved during anodal tDCS, might improve learning processes. On the other hand, a pure modulation of synaptic strength prior to learning might compromise performance, due to homeostatic or defocusing effects. Therefore, administering tDCS during, and not before, motor learning might be the best strategy to improve the effects of physical therapy [126].

The parameters of stimulation – such as number of stimulation sessions, frequency, intensity and site of stimulation – need to be taken in consideration. Relative to the duration of the cortical stimulation, it is worth mentioning that NIBS interventions have relatively short-lived after-effects compared to experimental LTP/LTD or to the duration needed for any clinically relevant functional improvement. However, repeated sessions of NIBS may have cumulative effects; perhaps due to these cumulative effects, several sessions of NIBS are usually associated with greater magnitude and duration of behavioural effects [129]. This has been also reported in clinical trials in stroke patients, in which stimulation with rTMS for 10 days can indeed induce a long-lasting improvement of motor behaviour that lasted for 10 days after the end of stimulation [32,112]; similarly, cathodal tDCS applied over 5 consecutive days is associated with a cumulative motor function improvement that lasts up to 2 weeks after the end of stimulation. However, interesting, this effect is not observed when sessions are applied weekly instead of daily [73]. In fact, multiple stimulation sessions are required in order to induce a significant manipulation in synaptic efficacy [130]. Thus, future clinical trails need to take into account that only prolonged and consecutive sessions of NIBS can translate into a long-lasting functional gains in stroke patients.

Until now physical therapy has been largely combined with NIBS applied to the motor cortex; nonetheless, other brain areas might be involved in motor recovery. For instance, higher levels of contralesional activity in pre-

frontal and parietal cortices appear to be predictive of a slower motor recovery, suggesting a possible negative role of activity in these areas of the intact hemisphere in functional restoration [38,131]. If so, suppression of such activity with NIBS might be a valuable intervention. Thus, modulation of excitability in areas beyond the primary motor cortex should be also taken into account as potentially interacting with the damaged motor areas, driving their activity-dependent activation. In patients with TBI, given their additional attentional impairments which negatively impact the efficacy of standard motor therapies [132], a modulation of attentional networks might enhance the responsiveness to standard motor rehabilitation.

Another controversial issue is related to the side of stimulation. It is still unclear whether it is better to suppress activity in the undamaged hemisphere or increase activity in the perilesional cortex. To date only one study, using tDCS, has directly compared the effectiveness of down-regulating the contralesional hemisphere with facilitation of the stroke hemisphere in patients with motor stroke; both approaches were found to be equally effective, with slightly greater improvement after suppression of the intact hemisphere [71]. However this was a small study and patient selection might have played a significant role. It is not yet known whether this is also true for rTMS. At least it appears that application of excitatory rTMS protocols to the stroke hemisphere is safe and does not increase the risk of provoking a seizure [133]. In any case, it is likely that rather than a global modulation of one or another hemisphere, more targeted, focal modulation of activity in selected cortical regions of each hemisphere might be desirable. Furthermore, the application of different strategies in different phases following the brain insult might be needed. Finally, it is worth remembering that currents induced by NIBS in the lesioned brain can be perturbed by anatomical changes which can render the neuromodulatory effects less predictable [52].

Importantly, the effects of NIBS are also task dependent; therefore it is possible that some motor tasks are more susceptible to modulation by NIBS than others. If so, the choice of the motor training task might be a critical determinant for the success of the therapy.

Overall, if guided by a careful consideration of the underlying mechanisms, the combination of NIBS with functional therapies has the potential to drive plastic changes in brain-damaged patients. This might in turn promote remarkable clinical gains in motor functions that otherwise could not be achieved by administering NIBS or motor treatment alone. Clearly, further investigation is warranted to address the overall utility of NIBS as an adjunct to stroke rehabilitation, and the optimal strategy to

combine the two interventions in order to maximize their functional interaction.

Conclusion

The uninjured tissue may be particularly receptive to modulation by various external tools including behavioral training and neuromodulatory approaches such as noninvasive brain stimulation. Given that both strategies, motor learning and cortical stimulation, have some similarities in their mechanisms of action, such as both induce similar changes in the local excitability in the lesioned and contralesional motor cortical area associated with long-lasting after-effects, their combination might be more beneficial than their use alone. In fact, brain stimulation can prime cortical excitability for a subsequent motor training task therefore optimizing processes of motor learning involved in standard rehabilitation therapies, leading to more pronounced and longer lasting functional gains. Some preliminary evidence seems to support this view. However, other studies failed to demonstrate a significant effect of brain stimulation as an adjuvant to standard motor therapy. In the future, the successful implementation of combined NIBS and motor therapy will critically rely on improved understanding of their functional interactions and associated effects on neural plasticity. Greater understanding of the mechanisms of action of each approach is necessary in order to optimize their combined use in rehabilitation and realize the promise of a more effective means to promote functional recovery after brain injury.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

NB, APL and FF conceived the initial idea. NB and FF wrote the first draft and all authors revised and approved the final manuscript.

References

- Desrosiers J, Bourbonnais D, Corriveau H, Gosselin S, Bravo G: **Effectiveness of unilateral and symmetrical bilateral task training for arm during the subacute phase after stroke: a randomized controlled trial.** *Clin Rehabil* 2005, **19**:581-593.
- Gillot AJ, Holder-Walls A, Kurtz JR, Varley NC: **Perceptions and experiences of two survivors of stroke who participated in constraint-induced movement therapy home programs.** *Am J Occup Ther* 2003, **57**:168-176.
- Luke C, Dodd KJ, Brock K: **Outcomes of the Bobath concept on upper limb recovery following stroke.** *Clin Rehabil* 2004, **18**:888-898.
- Roth EJ, Heinemann AW, Lovell LL, Harvey RL, McGuire JR, Diaz S: **Impairment and disability: their relation during stroke rehabilitation.** *Arch Phys Med Rehabil* 1998, **79**:329-335.
- Urton ML, Kohia M, Davis J, Neill MR: **Systematic literature review of treatment interventions for upper extremity hemiparesis following stroke.** *Occup Ther Int* 2007, **14**:11-27.
- Marshall S, Teasell R, Bayona N, Lippert C, Chundamala J, Villamere J, Mackie D, Cullen N, Bayley M: **Motor impairment rehabilitation post acquired brain injury.** *Brain Inj* 2007, **21**:133-160.
- Nelles G: **Cortical reorganization – effects of intensive therapy.** *Restor Neural Neurosci* 2004, **22**:239-244.
- Pascual-Leone A, Amedi A, Fregni F, Merabet LB: **The plastic human brain cortex.** *Annu Rev Neurosci* 2005, **28**:377-401.
- Nudo RJ, Friel KM: **Cortical plasticity after stroke: implications for rehabilitation.** *Rev Neurol (Paris)* 1999, **155**:713-717.
- Nudo RJ: **Plasticity.** *NeuroRx* 2006, **3**:420-427.
- Ward NS, Cohen LG: **Mechanisms underlying recovery of motor function after stroke.** *Arch Neurol* 2004, **61**:1844-1848.
- Hummelshheim H, Hauptmann B, Neumann S: **Influence of physiotherapeutic facilitation techniques on motor evoked potentials in centrally paretic hand extensor muscles.** *Electroencephalogr Clin Neurophysiol* 1995, **97**:18-28.
- Carey JR, Kimberley TJ, Lewis SM, Auerbach EJ, Dorsey L, Rundquist P, Ugurbil K: **Analysis of fMRI and finger tracking training in subjects with chronic stroke.** *Brain* 2002, **125**:773-788.
- Muellbacher W, Richards C, Ziemann U, Wittenberg G, Weltz D, Boroojerdi B, Cohen L, Hallett M: **Improving hand function in chronic stroke.** *Arch Neurol* 2002, **59**:1278-1282.
- Mark VW, Taub E: **Constraint-induced movement therapy for chronic stroke hemiparesis and other disabilities.** *Restor Neural Neurosci* 2004, **22**:317-336.
- Mudie MH, Matyas TA: **Can simultaneous bilateral movement involve the undamaged hemisphere in reconstruction of neural networks damaged by stroke?** *Disabil Rehabil* 2000, **22**:23-37.
- Mudie MH, Matyas TA: **Responses of the densely hemiplegic upper extremity to bilateral training.** *Neurorehabil Neural Repair* 2001, **15**:129-140.
- Buccino G, Solodkin A, Small SL: **Functions of the mirror neuron system: implications for neurorehabilitation.** *Cogn Behav Neurol* 2006, **19**:55-63.
- Ertelt D, Small S, Solodkin A, Dettmers C, McNamara A, Binkofski F, Buccino G: **Action observation has a positive impact on rehabilitation of motor deficits after stroke.** *Neuroimage* 2007, **36**(Suppl 2):T164-173.
- Fasoli SE, Krebs HI, Stein J, Frontera WR, Hogan N: **Effects of robotic therapy on motor impairment and recovery in chronic stroke.** *Arch Phys Med Rehabil* 2003, **84**:477-482.
- Fasoli SE, Krebs HI, Stein J, Frontera WR, Hughes R, Hogan N: **Robotic therapy for chronic motor impairments after stroke: Follow-up results.** *Arch Phys Med Rehabil* 2004, **85**:1106-1111.
- Bolton DA, Cauraugh JH, Hausenblas HA: **Electromyogram-triggered neuromuscular stimulation and stroke motor recovery of arm/hand functions: a meta-analysis.** *J Neurol Sci* 2004, **223**:121-127.
- Butler AJ, Page SJ: **Mental practice with motor imagery: evidence for motor recovery and cortical reorganization after stroke.** *Arch Phys Med Rehabil* 2006, **87**:S2-11.
- Schaechter JD: **Motor rehabilitation and brain plasticity after hemiparetic stroke.** *Prog Neurobiol* 2004, **73**:61-72.
- Hendricks HT, van Limbeek J, Geurts AC, Zwarts MJ: **Motor recovery after stroke: a systematic review of the literature.** *Arch Phys Med Rehabil* 2002, **83**:1629-1637.
- Fregni F, Pascual-Leone A: **Technology insight: noninvasive brain stimulation in neurology-perspectives on the therapeutic potential of rTMS and tDCS.** *Nat Clin Pract Neurol* 2007, **3**:383-393.
- Pascual-Leone A, Davey N, Rothwell JC, Wassermann E, B P: *Handbook of Transcranial Magnetic Stimulation* New York: Oxford University Press; 2002.
- Nitsche MA, Doemkes S, Karakose T, Antal A, Liebetanz D, Lang N, Tergau F, Paulus WW: **Shaping the effects of transcranial direct current stimulation of the human motor cortex.** *J Neurophysiol* 2007, **97**:3109-3117.
- Nitsche MA, Fricke K, Henschke U, Schlitterlau A, Liebetanz D, Lang N, Henning S, Tergau F, Paulus WW: **Pharmacological modulation of cortical excitability shifts induced by transcranial direct current stimulation in humans.** *J Physiol* 2003, **553**:293-301.
- Gandiga PC, Hummel FC, Cohen LG: **Transcranial DC stimulation (tDCS): a tool for double-blind sham-controlled clinical studies in brain stimulation.** *Clin Neurophysiol* 2006, **117**:845-850.
- Wagner T, Valero-Cabre A, Pascual-Leone A: **Noninvasive human brain stimulation.** *Annu Rev Biomed Eng* 2007, **9**:527-565.

32. Fregni F, Boggio PS, Valle AC, Rocha RR, Duarte J, Ferreira MJ, Wagner T, Fecteau S, Rigonatti SP, Riberto M, et al.: **A sham-controlled trial of a 5-day course of repetitive transcranial magnetic stimulation of the unaffected hemisphere in stroke patients.** *Stroke* 2006, **37**:2115-2122.
33. Fregni F, Boggio PS, Valle AC, Otachi P, Thut G, Rigonatti SP, Marcolin MA, Fecteau S, Pascual-Leone A, Fiore L, Valente K: **Homeostatic effects of plasma valproate levels on corticospinal excitability changes induced by 1 Hz rTMS in patients with juvenile myoclonic epilepsy.** *Clin Neurophysiol* 2006, **117**:1217-1227.
34. Fregni F, Santos CM, Myczkowski ML, Rigolino R, Gallucci-Neto J, Barbosa ER, Valente KD, Pascual-Leone A, Marcolin MA: **Repetitive transcranial magnetic stimulation is as effective as fluoxetine in the treatment of depression in patients with Parkinson's disease.** *J Neurol Neurosurg Psychiatry* 2004, **75**:1171-1174.
35. Triggs WJ, McCoy KJ, Greer R, Rossi F, Bowers D, Kortenkamp S, Nadeau SE, Heilman KM, Goodman WK: **Effects of left frontal transcranial magnetic stimulation on depressed mood, cognition, and corticomotor threshold.** *Biol Psychiatry* 1999, **45**:1440-1446.
36. Rossi S, Rossini PM: **TMS in cognitive plasticity and the potential for rehabilitation.** *Trends Cogn Sci* 2004, **8**:273-279.
37. Talelli P, Rothwell J: **Does brain stimulation after stroke have a future?** *Curr Opin Neurol* 2006, **19**:543-550.
38. Ward NS: **Mechanisms underlying recovery of motor function after stroke.** *Postgrad Med J* 2005, **81**:510-514.
39. Butefisch CM, Netz J, Wessling M, Seitz RJ, Homberg V: **Remote changes in cortical excitability after stroke.** *Brain* 2003, **126**:470-481.
40. Delvaux V, Alagona G, Gerard P, De Pasqua V, Pennisi G, de Noordhout AM: **Post-stroke reorganization of hand motor area: a 1-year prospective follow-up with focal transcranial magnetic stimulation.** *Clin Neurophysiol* 2003, **114**:1217-1225.
41. Shimizu T, Hosaki A, Hino T, Sato M, Komori T, Hirai S, Rossini PM: **Motor cortical disinhibition in the unaffected hemisphere after unilateral cortical stroke.** *Brain* 2002, **125**:1896-1907.
42. Murase N, Duque J, Mazzocchio R, Cohen LG: **Influence of inter-hemispheric interactions on motor function in chronic stroke.** *Ann Neurol* 2004, **55**:400-409.
43. Hummel FC, Cohen LG: **Non-invasive brain stimulation: a new strategy to improve neurorehabilitation after stroke?** *Lancet Neurol* 2006, **5**:708-712.
44. Barker AT, Jalinous R, Freeston IL: **Non-invasive magnetic stimulation of human motor cortex.** *Lancet* 1985, **1**:1106-1107.
45. Pascual-Leone A, Valls-Sole J, Wassermann EM, Hallett M: **Responses to rapid-rate transcranial magnetic stimulation of the human motor cortex.** *Brain* 1994, **117**(Pt 4):847-858.
46. Chen R, Classen J, Gerloff C, Celnik P, Wassermann EM, Hallett M, Cohen LG: **Depression of motor cortex excitability by low-frequency repetitive transcranial magnetic stimulation.** *Neurology* 1997, **48**:1398-1403.
47. Gangitano M, Valero-Cabre A, Tormos JM, Mottaghy FM, Romero JR, Pascual-Leone A: **Modulation of input-output curves by low and high frequency repetitive transcranial magnetic stimulation of the motor cortex.** *Clin Neurophysiol* 2002, **113**:1249-1257.
48. Maeda F, Keenan JP, Tormos JM, Topka H, Pascual-Leone A: **Inter-individual variability of the modulatory effects of repetitive transcranial magnetic stimulation on cortical excitability.** *Exp Brain Res* 2000, **133**:425-430.
49. Takeuchi N, Chuma T, Matsuo Y, Watanabe I, Ikoma KS: **Repetitive transcranial magnetic stimulation of contralesional primary motor cortex improves hand function after stroke.** *Stroke* 2005, **36**:2681-2686.
50. Talelli P, Greenwood RJ, Rothwell JC: **Exploring Theta Burst Stimulation as an intervention to improve motor recovery in chronic stroke.** *Clin Neurophysiol* 2007, **118**:333-342.
51. Di Lazzaro V, Pilato F, Dileone M, Profice P, Capone F, Ranieri F, Musumeci G, Cianfoni A, Pasqualetti P, Tonali PA: **Modulating cortical excitability in acute stroke: a repetitive TMS study.** *Clin Neurophysiol* 2008, **119**:715-723.
52. Wagner T, Fregni F, Eden U, Ramos-Estebanez C, Grodzinsky A, Zahn M, Pascual-Leone A: **Transcranial magnetic stimulation and stroke: a computer-based human model study.** *Neuroimage* 2006, **30**:857-870.
53. Hoffman RE, Cavus I: **Slow transcranial magnetic stimulation, long-term depotentiation, and brain hyperexcitability disorders.** *Am J Psychiatry* 2002, **159**:1093-1102.
54. Wang H, Wang X, Wetzel W, Scheich H: **Rapid-rate transcranial magnetic stimulation in auditory cortex induces LTP and LTD and impairs discrimination learning of frequency-modulated tones.** *Electroencephalogr Clin Neurophysiol Suppl* 1999, **51**:361-367.
55. Huang YZ, Edwards MJ, Rounis E, Bhatia KP, Rothwell JC: **Theta burst stimulation of the human motor cortex.** *Neuron* 2005, **45**:201-206.
56. Donoghue JP, Hess G, Sanes J: **Substrates and mechanisms for learning in motor cortex.** *Acquisition and Mechanisms for Learning in Motor Cortex* 1996:363-386.
57. Hess G, Donoghue JP: **Long-term depression of horizontal connections in rat motor cortex.** *Eur J Neurosci* 1996, **8**:658-665.
58. Hess G, Donoghue JP: **Long-term potentiation and long-term depression of horizontal connections in rat motor cortex.** *Acta Neurobiol Exp (Wars)* 1996, **56**:397-405.
59. Hasselmo ME: **Neuromodulation and cortical function: modeling the physiological basis of behavior.** *Behav Brain Res* 1995, **67**:1-27.
60. Strafella AP, Paus T, Fraraccio M, Dagher A: **Striatal dopamine release induced by repetitive transcranial magnetic stimulation of the human motor cortex.** *Brain* 2003, **126**:2609-2615.
61. Strafella AP, Paus T, Barrett J, Dagher A: **Repetitive transcranial magnetic stimulation of the human prefrontal cortex induces dopamine release in the caudate nucleus.** *J Neurosci* 2001, **21**:RC157.
62. Fujiki M, Steward O: **High frequency transcranial magnetic stimulation mimics the effects of ECS in upregulating astroglial gene expression in the murine CNS.** *Brain Res Mol Brain Res* 1997, **44**:301-308.
63. Hausmann A, Weis C, Marksteiner J, Hinterhuber H, Humpel C: **Chronic repetitive transcranial magnetic stimulation enhances c-fos in the parietal cortex and hippocampus.** *Brain Res Mol Brain Res* 2000, **76**:355-362.
64. Ji RR, Schlaepfer TE, Aizenman CD, Epstein CM, Qiu D, Huang JC, Rupp F: **Repetitive transcranial magnetic stimulation activates specific regions in rat brain.** *Proc Natl Acad Sci USA* 1998, **95**:15635-15640.
65. Muller MB, Toschi N, Kresse AE, Post A, Keck ME: **Long-term repetitive transcranial magnetic stimulation increases the expression of brain-derived neurotrophic factor and cholecystokinin mRNA, but not neuropeptide tyrosine mRNA in specific areas of rat brain.** *Neuropsychopharmacology* 2000, **23**:205-215.
66. Lisanby SH, Luber B, Perera T, Sackeim HA: **Transcranial magnetic stimulation: applications in basic neuroscience and neuropsychopharmacology.** *Int J Neuropsychopharmacol* 2000, **3**:259-273.
67. Brunoni A, Boggio P, Fregni F: **Can the 'yin and yang' BDNF hypothesis be used to predict the effects of rTMS treatment in neuropsychiatry?** *Med Hypotheses* 2008, **71**:279-282.
68. Paulus W: **Transcranial direct current stimulation (tDCS).** *Suppl Clin Neurophysiol* 2003, **56**:249-254.
69. Nitsche MA, Paulus W: **Sustained excitability elevations induced by transcranial DC motor cortex stimulation in humans.** *Neurology* 2001, **57**:1899-1901.
70. Romero JR, Ansel D, Sparing R, Gangitano M, Pascual-Leone A: **Subthreshold low frequency repetitive transcranial magnetic stimulation selectively decreases facilitation in the motor cortex.** *Clin Neurophysiol* 2002, **113**:101-107.
71. Fregni F, Boggio PS, Mansur CG, Wagner T, Ferreira MJ, Lima MC, Rigonatti SP, Marcolin MA, Freedman SD, Nitsche MA, Pascual-Leone A: **Transcranial direct current stimulation of the unaffected hemisphere in stroke patients.** *Neuroreport* 2005, **16**:1551-1555.
72. Hummel F, Cohen LG: **Improvement of motor function with noninvasive cortical stimulation in a patient with chronic stroke.** *Neurorehabil Neural Repair* 2005, **19**:14-19.
73. Boggio PS, Nunes A, Rigonatti SP, Nitsche MA, Pascual-Leone A, Fregni F: **Repeated sessions of noninvasive brain DC stimulation is associated with motor function improvement in stroke patients.** *Restor Neurol Neurosci* 2007, **25**:123-129.
74. Liebetanz D, Nitsche MA, Tergau F, Paulus W: **Pharmacological approach to the mechanisms of transcranial DC-stimula-**

- tion-induced after-effects of human motor cortex excitability. *Brain* 2002, **125**:2238-2247.
75. Nitsche MA, Jaussi W, Liebetanz D, Lang N, Tergau F, Paulus W: **Consolidation of human motor cortical neuroplasticity by D-cycloserine.** *Neuropsychopharmacology* 2004, **29**:1573-1578.
 76. Ardolino G, Bossi B, Barbieri S, Priori A: **Non-synaptic mechanisms underlie the after-effects of cathodal transcutaneous direct current stimulation of the human brain.** *J Physiol* 2005, **568**:653-663.
 77. Paulus W: **Outlasting excitability shifts induced by direct current stimulation of the human brain.** *Suppl Clin Neurophysiol* 2004, **57**:708-714.
 78. Wolters A, Sandbrink F, Schlottmann A, Kunesch E, Stefan K, Cohen LG, Benecke R, Classen J: **A temporally asymmetric Hebbian rule governing plasticity in the human motor cortex.** *J Neurophysiol* 2003, **89**:2339-2345.
 79. Cooper SJ: **Donald O. Hebb's synapse and learning rule: a history and commentary.** *Neurosci Biobehav Rev* 2005, **28**:851-874.
 80. Plautz EJ, Milliken GW, Nudo RJ: **Effects of repetitive motor training on movement representations in adult squirrel monkeys: role of use versus learning.** *Neurobiol Learn Mem* 2000, **74**:27-55.
 81. Zhuang P, Dang N, Waziri A, Gerloff C, Cohen LG, Hallett M: **Implicit and explicit learning in an auditory serial reaction time task.** *Acta Neurol Scand* 1998, **97**:131-137.
 82. Krakauer JW: **Motor learning: its relevance to stroke recovery and neurorehabilitation.** *Curr Opin Neurol* 2006, **19**:84-90.
 83. Pascual-Leone A, Nguyet D, Cohen LG, Brasil-Neto JP, Cammarota A, Hallett M: **Modulation of muscle responses evoked by transcranial magnetic stimulation during the acquisition of new fine motor skills.** *J Neurophysiol* 1995, **74**:1037-1045.
 84. Liepert J, Bauder H, Wolfgang HR, Miltner WH, Taub E, Weiller C: **Treatment-induced cortical reorganization after stroke in humans.** *Stroke* 2000, **31**:1210-1216.
 85. Chollet F, Weiller C: **Imaging recovery of function following brain injury.** *Curr Opin Neurobiol* 1994, **4**:226-230.
 86. Weiller C, Chollet F, Friston KJ, Wise RJ, Frackowiak RS: **Functional reorganization of the brain in recovery from striatocapsular infarction in man.** *Ann Neurol* 1992, **31**:463-472.
 87. Cramer SC, Nelles G, Benson RR, Kaplan JD, Parker RA, Kwong KK, Kennedy DN, Finklestein SP, Rosen BR: **A functional MRI study of subjects recovered from hemiparetic stroke.** *Stroke* 1997, **28**:2518-2527.
 88. Richards LG, Stewart KC, Woodbury ML, Senesac C, Cauraugh JH: **Movement-dependent stroke recovery: a systematic review and meta-analysis of TMS and fMRI evidence.** *Neuropsychologia* 2008, **46**:3-11.
 89. Lotze M, Braun C, Birbaumer N, Anders S, Cohen LG: **Motor learning elicited by voluntary drive.** *Brain* 2003, **126**:866-872.
 90. Karni A, Meyer G, Rey-Hipolito C, Jezzard P, Adams MM, Turner R, Ungerleider LG: **The acquisition of skilled motor performance: fast and slow experience-driven changes in primary motor cortex.** *Proc Natl Acad Sci USA* 1998, **95**:861-868.
 91. Classen J, Liepert J, Wise SP, Hallett M, Cohen LG: **Rapid plasticity of human cortical movement representation induced by practice.** *J Neurophysiol* 1998, **79**:1117-1123.
 92. Greenough WT, Larson JR, Withers GS: **Effects of unilateral and bilateral training in a reaching task on dendritic branching of neurons in the rat motor-sensory forelimb cortex.** *Behav Neural Biol* 1985, **44**:301-314.
 93. Van Reempts J, Dikova M, Werbrout L, Clincke G, Borgers M: **Synaptic plasticity in rat hippocampus associated with learning.** *Behav Brain Res* 1992, **51**:179-183.
 94. Neeper SA, Gomez-Pinilla F, Choi J, Cotman C: **Exercise and brain neurotrophins.** *Nature* 1995, **373**:109.
 95. Goelet P, Castellucci VF, Schacher S, Kandel ER: **The long and the short of long-term memory – a molecular framework.** *Nature* 1986, **322**:419-422.
 96. Rioult-Pedotti MS, Friedman D, Hess G, Donoghue JP: **Strengthening of horizontal cortical connections following skill learning.** *Nat Neurosci* 1998, **1**:230-234.
 97. Butefisch CM, Davis BC, Wise SP, Sawaki L, Kopylev L, Classen J, Cohen LG: **Mechanisms of use-dependent plasticity in the human motor cortex.** *Proc Natl Acad Sci USA* 2000, **97**:3661-3665.
 98. Grutzendler J, Kasthuri N, Gan WB: **Long-term dendritic spine stability in the adult cortex.** *Nature* 2002, **420**:812-816.
 99. Bury SD, Jones TA: **Unilateral sensorimotor cortex lesions in adult rats facilitate motor skill learning with the "unaffected" forelimb and training-induced dendritic structural plasticity in the motor cortex.** *J Neurosci* 2002, **22**:8597-8606.
 100. Campanac E, Debanne D: **Plasticity of neuronal excitability: Hebbian rules beyond the synapse.** *Arch Ital Biol* 2007, **145**:277-287.
 101. Fregni F, Liebetanz D, Monte-Silva KK, Oliveira MB, Santos AA, Nitsche MA, Pascual-Leone A, Guedes RC: **Effects of transcranial direct current stimulation coupled with repetitive electrical stimulation on cortical spreading depression.** *Exp Neurol* 2007, **204**:462-466.
 102. Liebetanz D, Fregni F, Monte-Silva KK, Oliveira MB, Amancio-dos-Santos A, Nitsche MA, Guedes RC: **After-effects of transcranial direct current stimulation (tDCS) on cortical spreading depression.** *Neurosci Lett* 2006, **398**:85-90.
 103. Nitsche MA, Roth A, Kuo MF, Fischer AK, Liebetanz D, Lang N, Tergau F, Paulus W: **Timing-dependent modulation of associative plasticity by general network excitability in the human motor cortex.** *J Neurosci* 2007, **27**:3807-3812.
 104. Siebner HR, Lang N, Rizzo V, Nitsche MA, Paulus W, Lemon RN, Rothwell JC: **Preconditioning of low-frequency repetitive transcranial magnetic stimulation with transcranial direct current stimulation: evidence for homeostatic plasticity in the human motor cortex.** *J Neurosci* 2004, **24**:3379-3385.
 105. Butefisch CM, Khurana V, Kopylev L, Cohen LG: **Enhancing encoding of a motor memory in the primary motor cortex by cortical stimulation.** *J Neurophysiol* 2004, **91**:2110-2116.
 106. Carey JR, Fregni F, Pascual-Leone A: **rTMS combined with motor learning training in healthy subjects.** *Restor Neurol Neurosci* 2006, **24**:191-199.
 107. Antal A, Nitsche MA, Kincses TZ, Kruse W, Hoffmann KP, Paulus W: **Facilitation of visuo-motor learning by transcranial direct current stimulation of the motor and extrastriate visual areas in humans.** *Eur J Neurosci* 2004, **19**:2888-2892.
 108. Liebetanz D, Nitsche MA, Paulus W: **Pharmacology of transcranial direct current stimulation: missing effect of riluzole.** *Suppl Clin Neurophysiol* 2003, **56**:282-287.
 109. Floel A, Rosser N, Michka O, Knecht S, Breitenstein C: **Noninvasive brain stimulation improves language learning.** *J Cogn Neurosci* 2008, **20**:1415-1422.
 110. Adkins-Muir DL, Jones TA: **Cortical electrical stimulation combined with rehabilitative training: enhanced functional recovery and dendritic plasticity following focal cortical ischemia in rats.** *Neurol Res* 2003, **25**:780-788.
 111. Brown JA, Lutsep HL, Weinand M, Cramer SC: **Motor cortex stimulation for the enhancement of recovery from stroke: a prospective, multicenter safety study.** *Neurosurgery* 2006, **58**:464-473.
 112. Khedr EM, Ahmed MA, Fathy N, Rothwell JC: **Therapeutic trial of repetitive transcranial magnetic stimulation after acute ischemic stroke.** *Neurology* 2005, **65**:466-468.
 113. Kim YH, You SH, Ko MH, Park JW, Lee KH, Jang SH, Yoo WK, Hallett M: **Repetitive transcranial magnetic stimulation-induced corticomotor excitability and associated motor skill acquisition in chronic stroke.** *Stroke* 2006, **37**:1471-1476.
 114. Takeuchi N, Tada T, Toshima M, Chuma T, Matsuo Y, Ikoma K: **Inhibition of the unaffected motor cortex by 1 Hz repetitive transcranial magnetic stimulation enhances motor performance and training effect of the paretic hand in patients with chronic stroke.** *J Rehabil Med* 2008, **40**:298-303.
 115. Pomeroy VM, Cloud G, Tallis RC, Donaldson C, Nayak V, Miller S: **Transcranial magnetic stimulation and muscle contraction to enhance stroke recovery: a randomized proof-of-principle and feasibility investigation.** *Neurorehabil Neural Repair* 2007, **21**:509-517.
 116. Malcolm MP, Triggs WJ, Light KE, Gonzalez Rothi LJ, Wu S, Reid K, Nadeau SE: **Repetitive transcranial magnetic stimulation as an adjunct to constraint-induced therapy: an exploratory randomized controlled trial.** *Am J Phys Med Rehabil* 2007, **86**:707-715.
 117. Wolf SL, Thompson PA, Morris DM, Rose DK, Winstein CJ, Taub E, Giuliani C, Pearson SL: **The EXCITE trial: attributes of the Wolf Motor Function Test in patients with subacute stroke.** *Neurorehabil Neural Repair* 2005, **19**:194-205.

118. Uswatte G, Taub E, Morris D, Vignolo M, McCulloch K: **Reliability and validity of the upper-extremity Motor Activity Log-14 for measuring real-world arm use.** *Stroke* 2005, **36**:2493-2496.
119. Desrosiers J, Bravo G, Hebert R, Dutil E, Mercier L: **Validation of the Box and Block Test as a measure of dexterity of elderly people: reliability, validity, and norms studies.** *Arch Phys Med Rehabil* 1994, **75**:751-755.
120. Fregni F, Gimenes R, Valle AC, Ferreira MJ, Rocha RR, Natale L, Bravo R, Rigonatti SP, Freedman SD, Nitsche MA, et al.: **A randomized, sham-controlled, proof of principle study of transcranial direct current stimulation for the treatment of pain in fibromyalgia.** *Arthritis Rheum* 2006, **54**:3988-3998.
121. Jebson RH, Taylor N, Trieschmann RB, Trotter MJ, Howard LA: **An objective and standardized test of hand function.** *Arch Phys Med Rehabil* 1969, **50**:311-319.
122. Hesse S, Werner C, Schonhardt EM, Bardeleben A, Jenrich W, Kirker SG: **Combined transcranial direct current stimulation and robot-assisted arm training in subacute stroke patients: a pilot study.** *Restor Neurol Neurosci* 2007, **25**:9-15.
123. Thickbroom GW: **Transcranial magnetic stimulation and synaptic plasticity: experimental framework and human models.** *Exp Brain Res* 2007, **180**:583-593.
124. Turrigiano GG, Nelson SB: **Homeostatic plasticity in the developing nervous system.** *Nat Rev Neurosci* 2004, **5**:97-107.
125. Hummel FC, Cohen LG: **Drivers of brain plasticity.** *Curr Opin Neurol* 2005, **18**:667-674.
126. Kuo MF, Unger M, Liebetanz D, Lang N, Tergau F, Paulus W, Nitsche MA: **Limited impact of homeostatic plasticity on motor learning in humans.** *Neuropsychologia* 2008, **46**:2122-2128.
127. Nitsche MA, Schauenburg A, Lang N, Liebetanz D, Exner C, Paulus W, Tergau F: **Facilitation of implicit motor learning by weak transcranial direct current stimulation of the primary motor cortex in the human.** *J Cogn Neurosci* 2003, **15**:619-626.
128. Canepari M, Djuricic M, Zecevic D: **Dendritic signals from rat hippocampal CA1 pyramidal neurons during coincident pre- and post-synaptic activity: a combined voltage- and calcium-imaging study.** *J Physiol* 2007, **580**:463-484.
129. Gershon AA, Dannon PN, Grunhaus L: **Transcranial magnetic stimulation in the treatment of depression.** *Am J Psychiatry* 2003, **160**:835-845.
130. Froc DJ, Chapman CA, Trepel C, Racine RJ: **Long-term depression and depotentiation in the sensorimotor cortex of the freely moving rat.** *J Neurosci* 2000, **20**:438-445.
131. Webster BR, Celnik PA, Cohen LG: **Noninvasive brain stimulation in stroke rehabilitation.** *NeuroRx* 2006, **3**:474-481.
132. Ziino C, Ponsford J: **Selective attention deficits and subjective fatigue following traumatic brain injury.** *Neuropsychology* 2006, **20**:383-390.
133. Yozbatiran N, Alonso-Alonso M, See J, Demirtas-Tatlidede A, Luu D, Motiwala RR, Pascual-Leone A, Cramer SC: **Safety and behavioral effects of high-frequency repetitive transcranial magnetic stimulation in stroke.** *Stroke* 2009, **40**:309-312.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

