



NIH Public Access

Author Manuscript

Brain Stimul. Author manuscript; available in PMC 2009 October 1.

Published in final edited form as:
Brain Stimul. 2008 October ; 1(4): 363–369. doi:10.1016/j.brs.2008.08.001.

Consensus: “Can tDCS and TMS enhance motor learning and memory formation?”

Janine Reis¹, Edwin Robertson², John W. Krakauer³, John Rothwell⁴, Lisa Marshall⁵, Christian Gerloff⁶, Eric Wassermann⁷, Alvaro Pascual-Leone², Friedhelm Hummel⁶, Pablo A. Celnik⁸, Joseph Classen⁹, Agnes Floel¹⁰, Ulf Ziemann¹¹, Walter Paulus¹², Hartwig R. Siebner¹³, Jan Born⁵, and Leonardo G. Cohen¹

¹ Human Cortical Physiology Section, NINDS, NIH, Bethesda, MD, USA

² Berenson-Allen Center for Noninvasive Brain Stimulation, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA

³ Motor Control Laboratory, Dept. of Neurology, Columbia University, New York, USA

⁴ Institute of Neurology, Queen Square, London, UK

⁵ Department of Neurology and Clinical Neuroendocrinology, University of Lübeck, Germany

⁶ Department of Neurology, University of Hamburg, Germany

⁷ Brain Stimulation Unit, NINDS, NIH, Bethesda, MD, USA

⁸ Dept. of Physical Medicine and Rehabilitation, Johns Hopkins University, Baltimore, MD, USA

⁹ Human Cortical Physiology and Motor Control Laboratory, Department of Neurology, University of Würzburg, Germany

¹⁰ Department of Neurology, University of Münster and IZKF Münster, Germany

¹¹ Department of Neurology, Goethe-University of Frankfurt, Germany

¹² Department of Clinical Neurophysiology, University of Goettingen, Germany

¹³ Brain mapping laboratory, Department of Neurology, Christian-Albrechts University, Kiel, Germany

Abstract

Noninvasive brain stimulation has developed as a promising tool for cognitive neuroscientists. Transcranial magnetic (TMS) and direct current (tDCS) stimulation allow researchers to purposefully enhance or decrease excitability in focal areas of the brain. The purpose of this paper is to review information on the use of TMS and tDCS as research tools to facilitate motor memory formation, motor performance and motor learning in healthy volunteers. Studies implemented so far have mostly focused on the ability of TMS and tDCS to elicit relatively short lasting motor improvements and the mechanisms underlying these changes have been only partially investigated. Despite limitations including the scarcity of data, work that has been already accomplished raises the exciting hypothesis

Proofs and correspondence to: Leonardo G. Cohen, Human Cortical Physiology Section, National Institute of Neurological Disorders and Stroke, National Institutes of Health, 10 Center Drive, Bethesda, MD 20892-1430, USA, Phone: (301) 496-9782, Fax: (301) 480-2286, E-mail: cohenl@ninds.nih.gov.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

that currently available noninvasive transcranial stimulation techniques could modulate motor learning and memory formation in healthy humans and potentially in patients with neurological and psychiatric disorders.

Introduction

Within the past two decades noninvasive brain stimulation has been used as a probe to modulate attention, memory, motor and language functions in humans.^{1–9} TMS and tDCS can enhance or decrease excitability in target cortical regions depending on the parameters of stimulation utilized.^{10–13} TMS, and to a lesser extent tDCS (please see Chapter by Nitsche et al. for specific differences between tDCS and TMS), have been used as an interference technique (“virtual lesion”) for understanding brain-behavior interactions and to explore possible cause-effect links between altered activity in specific brain areas and particular behaviors.^{14, 15} Improved understanding of the involvement of a brain region in a type of behavior was followed by attempts to modify activity in this area to secondarily influence performance, learning and memory functions.^{2–9} In this chapter we summarize results from studies aiming at eliciting improvements in motor performance and motor learning in healthy humans.

Motor learning

Formation of motor memories is required for learning of motor skills in daily life.^{16, 17} It is helpful to distinguish studies focused on the process of acquisition, consolidation and long-term stability (also referred to as retention) of a new motor skill^{8, 18, 19} from those that evaluate return to baseline levels of performance in response to external perturbations.^{20–23} An example of adaptation to an external perturbation is the response to directional errors in visually-guided reaching movement caused by prism glasses^{22, 24}: with practice, performance returns to the “baseline” level. Importantly, adaptation may not require acquisition of new motor synergies or movement patterns, as it engages movements that were achieved throughout life.

In contrast to adaptation, acquisition of a new motor skill involves the acquisition of new movement qualities and/or muscle synergies that enhance performance beyond pre-existing levels. Skills seem to take longer to acquire than adaptation and sometimes do not reach plateau levels after years (i.e., learning to play piano or basketball).^{20, 25–27} In engineering terms, adaptation may be modeled as error based learning whereas, for example, motor skill learning is better modeled in terms of reward-based signals.^{28, 29} However, it should be kept in mind that an overlap of reward and error based learning is possible.

Consolidation refers either to stabilization (reduced susceptibility to retrograde interference) or off-line improvements.^{30–33} Consolidation processes can depend on the type of task, the time between end of practice and testing of recall, and the presence or absence of sleep. Off-line improvements, for instance in the ability to perform a finger opposition task, correlated with the amount of time spent in REM sleep.³⁴ Similarly, off-line improvements in a motor sequence learning task are sleep-dependent when individuals are aware of the underlying sequence.³⁵ However, when individuals have little awareness for the sequence, off-line improvements are able to develop over waking or over a night sleep.³⁵ Potentially, the effect of individuals’ awareness upon off-line learning is mediated by their declarative knowledge for the sequence: disrupting declarative knowledge for the sequence can induce improvements over wake.³⁶ In adaptation studies, the successful return to baseline performance after the perturbation occurs often within one session, and therefore the possibility of offline improvements across days has not been thoroughly tested, although savings, an increase in the rate of re-adaptation, could be considered a form of off-line learning.³⁷ One exception to this general statement is the study of Huber et al. in which overnight improvements in performance

of a motor adaptation paradigm in which subjects had to adapt to a visual perturbation of a reaching movement correlated with increased <4 Hz activity during slow wave sleep.³⁸ This activity is thought to reflect oscillatory changes in neuronal membrane potentials.³⁸

As stated above, another form of consolidation is stabilization, that is, maintenance of practice-induced performance improvements or skill (in opposition to forgetting or to off-line improvements). After the end of a practice period, procedural memories for a task A may display different degrees of strength to interference. A classical approach to evaluate this strength is to introduce a task B as a source of interference and subsequently test the subject's ability to perform task A.^{37, 39, 40} Interference to recall a newly learned motor task A by practicing a different motor task B has been described as "retroactive interference". Retroactive interference has a well described time-course diminishing with the length of the time interval between the end of practice of task A and the application of the interfering task B, becoming virtually absent after 6 hours.^{37, 40, 41} Stabilization over hours following learning of dynamic adaptation tasks has been well documented.^{41, 42} Of note, Goedert and Willingham showed that for motor sequence learning off-line stabilization does not occur. Whether 15 minutes or 24 hours passes in between task A and B, task B continues to exert retrograde interference.⁴⁰ However, this finding might be task specific.

It is important to keep in mind that during motor learning, both kinds of consolidation, off-line learning and stabilization, are likely occurring.^{43, 44} As stated above, it is unclear if off-line enhancement occurs for adaptation tasks as consolidation studies of adaptation have focused on stabilization. Other factors that may influence our ability to assess the stability of a procedural memory include the endpoint measure utilized (speed, accuracy, or speed accuracy ratios related to a motor task) and the practice schedule^{45, 46}: for instance, when a skill is acquired through interleaved rather than blocked practice schedules, motor memories may become more resistant to interference.

One important theoretical point to highlight is the difference between measurements of motor performance and motor skill. Improvements in speed or in accuracy of performance of a motor action have been often reported in isolation in the literature, occasionally indicating that changes in one of these two measures occurred in the absence of changes in the other. Such changes have been reported as changes in skill. It would be important to keep in mind that motor skill cannot always be reliably surmised from changes in only one of these two measures. Skill may be better described as a change in the speed-accuracy trade-off, which is task dependent. Taking into consideration this issue, would help future investigators avoid concluding a change in skill (skill improvements) when in fact subjects have only moved along the same speed/accuracy trade-off curve.

Non-invasive brain stimulation

Non-invasive brain stimulation has been used to identify the functional relevance of particular brain regions in motor learning and facilitate activity in specific cortical areas involved in motor learning in an attempt to improve motor function.

a. Functional role of the primary motor cortex in motor learning as studied with noninvasive cortical stimulation

Motor learning is associated with functional changes in a distributed network that includes the primary motor, premotor and supplementary motor cortices, the cerebellum, thalamic nuclei and the striatum^{18, 47–50}. Most TMS and tDCS studies carried out so far to study the role of motor areas in motor learning have focused on M1.

Role of M1 in encoding of an elementary motor memory—Butefisch et al. showed that the synchronous application of single pulse TMS to M1 contralateral to a hand practicing a thumb abduction task⁵¹ enhanced the ability of healthy subjects to encode an elementary and short-lasting motor memory in the primary motor cortex.² Importantly, this effect was evident when M1 was stimulated in synchrony with the training motions but not when applied in between training movements. A second important finding from this study was that synchronous stimulation of the “resting” M1 with the training motions in the ipsilateral hand, cancelled training effects on motor memory formation, consistent with the hypothesis that interhemispheric interactions between M1s contribute to motor memory formation.² It has been proposed that formation of motor memories within M1 could represent a first step in the more complex chain of events leading to improve a motor skill, but it should not be interpreted as motor learning per se as skill improvements above naïve levels are typically not seen with this particular paradigm of motor memory formation.

Role of M1 in motor adaptation—As discussed above, the concept of motor adaptation refers to learning to adjust to external perturbations.^{21–23, 52} In these experiments, subjects adjust their motor behavior to compensate for a particular perturbation in order to maintain a stable performance.²¹ TMS and tDCS have been used to evaluate the role of M1 in motor adaptation. In general, stimulation over M1, using parameters that decrease excitability in that region like 1Hz TMS, have been applied before or during adaptation paradigms (in order to evaluate its functional relevance for encoding of the necessary adjustments to compensate for the perturbation).^{23, 53} In one study, single TMS pulses applied to M1 at 120% of RMT of the first dorsal interosseus muscle immediately after the end of each trial while adapting to a perturbation in the form of a visuomotor rotation did not impact adaptation, but caused faster de-adaptation (forgetting) within the same session relative to single pulses applied 700 ms after the end of each trial or relative to PMd stimulation.⁵³ In another study, 1 Hz rTMS applied to M1 at 90 % of biceps RMT before force field adaptation did not affect the participants’ adaptation per se, but impaired retention relative to control subjects (who did not receive any rTMS) as tested the following day, when subjects were exposed to the same force field to which they had previously adapted.⁵⁴ On the other hand, Baraduc et al. did not find a deleterious effect of 1Hz TMS applied over M1 on adaptation to a dynamic force field.²³ Potential areas of interest that remain to be investigated in more detail include the role of motor areas other than M1 in motor adaptation.

Role of M1 in motor skill learning—As discussed above, motor learning may (and often does) continue after the end of practice periods, referred to above as consolidation in the form of off-line learning. The role of M1 has been investigated in the process of acquisition and consolidation of motor skills.^{13, 50–55} Muellbacher et al. reported in an influential study, that 1 Hz rTMS over M1 at 115% of flexor pollicis brevis RMT applied immediately before a single session practice of a thumb-to-finger opposition task did not disrupt within session improvements in speed and muscle force generation but had deleterious effects on retention of these improvements as tested the following day relative to stimulation applied 6 hr after practice or when applied to other cortical areas such as the occipital cortex or the left DLPFC.¹³ These results were interpreted as supportive of the view that M1 plays a functionally relevant role in consolidation of explicit motor memories.¹³ The role of M1 was also explored in motor sequence learning⁵⁶ using a modified version of the serial reaction time task (SRTT).⁵⁷ 1 Hz rTMS was applied over M1 immediately after training when subjects practiced the task early in the morning or late in the evening (different groups). The endpoint measure was off-line enhancement in performance of the task 12 hours later (the evening of the practice day with no sleep in between in the first group and the following morning after a normal night sleep in the second group). It was reported that off-line enhancements of the learned task were disrupted in the first group (no sleep) but not in the second group. The authors interpreted the result as

indicative of different consolidation processes depending on how close sleep is to the practice period.⁵⁶ For a more detailed review of the effects of virtual lesion studies on motor learning, please refer to¹⁵. On the other hand, rTMS and tDCS have been used in an attempt to facilitate motor learning, the focus of this review.

b. Attempts to facilitate motor learning by noninvasive cortical stimulation

Based on human neuroimaging studies, it was proposed that application of noninvasive stimulation with parameters that enhance motor cortical excitability could secondarily facilitate motor learning. One key structure in the distributed network engaged in motor sequence learning is the primary motor cortex (M1).^{13, 58} Within M1 the extent of cortical reorganization associated with motor training correlates with performance improvements.⁵⁹ The interaction between the two M1s appears to play an important role in motor control in general^{20, 60–64} and in motor sequence learning in particular.^{64, 65} However, the specific way in which these interactions operate during motor learning remain to be determined. According to these interactions, it would be theoretically possible to facilitate motor learning processes in which M1 is involved by enhancing excitability in the “learning” M1⁵ or by decreasing excitability in the “resting” M1^{66, 67}, but see also Wassermann et al.⁶⁸ The intrinsic intracortical mechanisms by which these oversimplified models may operate remain to be identified^{64, 69, 70} (see for discussion chapters by Walsh et al., Di Lazzaro et al., Berardelli et al.).

Several investigators proposed that noninvasive cortical stimulation that enhances excitability in the M1 contralateral to a training hand^{5, 71} might result in varying degrees of improvement in motor function in healthy humans. Kim et al reported that 10 Hz rTMS at an intensity of 80% of resting motor threshold (RMT) applied over M1 during practice of contralateral sequential finger movements resulted in improved motor sequence learning (as measured by target accuracy and speed on a sequential key press task) relative to sham stimulation.⁷² Anodal tDCS applied over M1 during practice also led to improvements in: (a) the number of correct key presses in a sequential finger movement task in a polarity-specific manner since cathodal tDCS failed to induce this effect⁷³, (b) performance of a visuomotor coordination task that were transient (~ 5 minutes)⁷⁴, (c) reaction times in the sequence blocks relative to the random blocks in the SRTT⁵⁷, in which subjects learn a sequence of 12 key presses without explicit awareness⁷⁵, (d) performance of the Jebsen Taylor Hand function test (JTT), a task often used in stroke research that mimics activities of daily living such as lifting cans and picking up small objects.⁷⁶ Interestingly, this effect has been reported as present in the nondominant hand only in young healthy adults.⁷⁷ Of note, tDCS in these two papers was applied after subjects reached stable JTT performance, likely reflecting a tDCS-induced performance improvement beyond a plateau level. It is not known if application of noninvasive cortical stimulation during the learning period of the task (before it reaches an asymptote) could speed up or enhance learning of the task.

In contrast to studies that focused on application of TMS or tDCS to the M1 contralateral to a practicing hand, the application of 1Hz rTMS to the M1 ipsilateral to a training hand results in: (a) increases in motor cortical excitability of the opposite M1^{66, 67, 78} relative to sham stimulation, and (b) improvements in motor sequence learning⁶⁵ relative to stimulation of the contralateral M1, ipsilateral premotor area, or vertex (Cz). One important consideration is that the effects of stimulating M1 with either TMS or tDCS are likely to be dependent on the complexity of the task. For example, performance of relatively simple repetitive finger abduction movements was not improved by high frequency rTMS over the “learning” M1⁷⁹ while more complex sequential motor tasks or encoding of a motor memory did improve (see above). Similarly, another study showed no performance improvement in a task engaging single finger tracking motions when the ipsilateral M1 was stimulated with 1Hz rTMS during practice.⁸⁰

In contrast to studies focusing on motor sequence learning or motor performance, we are not aware of studies that used TMS or tDCS in an attempt to facilitate motor adaptation.

Caveats and future directions

While the previous paragraphs depict a relatively consistent and homogenous picture on the effects of up- and down-regulation of excitability within M1 on motor learning and motor memory formation, several caveats should be kept in mind. **First**, induction of a “virtual lesion” or enhancement of activity in one cortical area may result in behavioral changes through specific effects on that area or secondarily through distant effects on other interconnected cortical areas. **Second**, the discussion of results above (as well as those of specific physiological interactions across cortical regions⁶¹) assume in general that the surface of the brain is a smooth sphere, often neglecting that the folding of the cortex may result in hyperpolarization of neurons on one side of a gyrus but depolarization on the other. **Third**, the history of activity in the stimulated cortex may be of considerable importance. For example, the effects of stimulation may differ substantially if applied to a “fatigued” or to a “well rested” cortex. In one example, 1 Hz rTMS may induce facilitatory effects if acting on a cortex that has been previously inhibited by cathodal tDCS.^{81, 82} This phenomenon, referred to as homeostatic plasticity or metaplasticity and discussed elsewhere⁸³, may potentially impact motor learning.^{84, 85} **Fourth**, while most work in the field focused on studying the effects of stimulation over M1, some reports indicated that stimulation of the dorsal premotor or the lateral or medial prefrontal cortex failed to induce overt reaction time improvements in the SRTT task.⁷⁵ Similarly, anodal tDCS applied over the primary visual cortex (V1) did not improve performance of a visuomotor tracking task in healthy subjects.⁷⁴ However, findings indicating a lack of effect of stimulation in a particular site is not proof that that area is not involved. The most parsimonious interpretation of these findings is that more elaborated, hypothesis-driven behavioral paradigms or stimulation strategies may be necessary to study the functional role of these cortical regions in motor learning and memory formation. Alternatively, it is possible that the “threshold” for facilitating motor learning by M1 stimulation is lower than by stimulation of other cortical areas, an issue to be investigated in future experiments. **Fifth**, one caveat of many previous investigations has been the focus on short term improvements in performance. More experiments are required to assess the effects of repeated applications of TMS or tDCS in association with multiple training sessions, their interaction with specific motor learning stages and tasks and the extent to which these performance improvements are retained in the long term. **Finally**, it should be kept in mind that the effects of TMS and tDCS can not be assumed to be the same. The mechanisms underlying the effects of each technique on motor cortical function are very likely to differ (see chapter by Nitsche et al.) and so will the effects on motor cortical networks and behavioral consequences of its application. Clearly more work is needed in this area.

Summary

In summary, the scarce studies performed so far point to the encouraging conclusion that noninvasive brain stimulation can contribute to the understanding of mechanisms underlying motor learning and motor memory formation and raise the exciting hypothesis that this increased understanding could in the future result in the development of new strategies to enhance specific stages of learning and memory processing in healthy humans and in patients with brain lesions (see chapter by Gerloff et al.).

References

1. Jahanshahi M, Rothwell J. Transcranial magnetic stimulation studies of cognition: an emerging field. *Exp Brain Res* 2000;131:1–9. [PubMed: 10759166]

2. Butefisch CM, Khurana V, Kopylev L, Cohen LG. Enhancing encoding of a motor memory in the primary motor cortex by cortical stimulation. *J Neurophysiol* 2004;91:2110–2116. [PubMed: 14711974]
3. Duque J, Mazzocchio R, Stefan K, et al. Memory Formation in the Motor Cortex Ipsilateral to a Training Hand. *Cereb Cortex*. 2007
4. Hummel F, Celnik P, Giroux P, et al. Effects of non-invasive cortical stimulation on skilled motor function in chronic stroke. *Brain*. 2005
5. Pascual-Leone A, Valls-Sole J, Wassermann EM, Hallett M. Responses to rapid-rate transcranial magnetic stimulation of the human motor cortex. *Brain* 1994;117(Pt 4):847–858. [PubMed: 7922470]
6. Wassermann EM. Risk and safety of repetitive transcranial magnetic stimulation: report and suggested guidelines from the International Workshop on the Safety of Repetitive Transcranial Magnetic Stimulation, June 5–7, 1996. *Electroencephalogr Clin Neurophysiol* 1998;108:1–16. [PubMed: 9474057]
7. Flöel A, Rösler N, Michka O, et al. Noninvasive Brain Stimulation Improves Language Learning. *J Cogn Neurosci*. 2008
8. Robertson E, Press D, Pascual-Leone A. Off-line learning and the primary motor cortex. *J Neurosci* 2005;25:6372–6378. [PubMed: 16000627]
9. Marshall L, Helgadóttir H, Mölle M, Born J. Boosting slow oscillations during sleep potentiates memory. *Nature* 2006;444:610–613. [PubMed: 17086200]
10. Chen R, Classen J, Gerloff C, et al. Depression of motor cortex excitability by low-frequency transcranial magnetic stimulation. *Neurology* 1997;48:1398–1403. [PubMed: 9153480]
11. Nitsche MA, Liebetanz D, Tergau F, Paulus W. [Modulation of cortical excitability by transcranial direct current stimulation]. *Nervenarzt* 2002;73:332–335. [PubMed: 12040980]
12. Bestmann S, Baudewig J, Siebner HR, et al. Subthreshold high-frequency TMS of human primary motor cortex modulates interconnected frontal motor areas as detected by interleaved fMRI-TMS. *Neuroimage* 2003;20:1685–1696. [PubMed: 14642478]
13. Muellbacher W, Ziemann U, Wissel J, et al. Early consolidation in human primary motor cortex. *Nature* 2002;415:640–644. [PubMed: 11807497]
14. Cohen LG, Celnik P, Pascual-Leone A, et al. Functional relevance of cross-modal plasticity in blind humans. *Nature* 1997;389:180–183. [PubMed: 9296495]
15. Siebner HR, Rothwell J. Transcranial magnetic stimulation: new insights into representational cortical plasticity. *Exp Brain Res* 2003;148:1–16. [PubMed: 12478392]
16. Squire LR. Memory systems of the brain: a brief history and current perspective. *Neurobiol Learn Mem* 2004;82:171–177. [PubMed: 15464402]
17. Dudai Y. The neurobiology of consolidations, or, how stable is the engram? *Annu Rev Psychol* 2004;55:51–86. [PubMed: 14744210]
18. Karni A, Meyer G, Rey-Hipolito C, et al. The acquisition of skilled motor performance: fast and slow experience-driven changes in primary motor cortex. *Proc Natl Acad Sci U S A* 1998;95:861–868. [PubMed: 9448252]
19. Walker MP, Brakefield T, Morgan A, et al. Practice with sleep makes perfect: sleep-dependent motor skill learning. *Neuron* 2002;35:205–211. [PubMed: 12123620]
20. Shadmehr R, Krakauer J. A computational neuroanatomy for motor control. *Exp Brain Res* 2008;185:359–381. [PubMed: 18251019]
21. Izawa J, Rane T, Donchin O, Shadmehr R. Motor adaptation as a process of reoptimization. *J Neurosci* 2008;28:2883–2891. [PubMed: 18337419]
22. Fernández-Ruiz J, Díaz R. Prism adaptation and aftereffect: specifying the properties of a procedural memory system. *Learn Mem* 1999;6:47–53. [PubMed: 10355523]
23. Baraduc P, Lang N, Rothwell JC, Wolpert DM. Consolidation of dynamic motor learning is not disrupted by rTMS of primary motor cortex. *Curr Biol* 2004;14:252–256. [PubMed: 14761660]
24. Kitazawa S, Kimura T, Uka T. Prism adaptation of reaching movements: specificity for the velocity of reaching. *J Neurosci* 1997;17:1481–1492. [PubMed: 9006989]

25. Hund-Georgiadis M, von Cramon DY. Motor-learning-related changes in piano players and non-musicians revealed by functional magnetic-resonance signals. *Exp Brain Res* 1999;125:417–425. [PubMed: 10323287]
26. Memmert D. Long-term effects of type of practice on the learning and transfer of a complex motor skill. *Percept Mot Skills* 2006;103:912–916. [PubMed: 17326522]
27. Karni A, Sagi D. The time course of learning a visual skill. *Nature* 1993;365:250–252. [PubMed: 8371779]
28. Shadmehr R, Krakauer JW. A computational neuroanatomy for motor control. *Exp Brain Res* 2008;185:359–381. [PubMed: 18251019]
29. Smith MA, Ghazizadeh A, Shadmehr R. Interacting adaptive processes with different timescales underlie short-term motor learning. *PLoS Biol* 2006;4:e179. [PubMed: 16700627]
30. Stickgold R. Sleep-dependent memory consolidation. *Nature* 2005;437:1272–1278. [PubMed: 16251952]
31. Walker M, Brakefield T, Seidman J, et al. Sleep and the time course of motor skill learning. *Learn Mem* 2003;10:275–284. [PubMed: 12888546]
32. Gais S, Albouy G, Boly M, et al. Sleep transforms the cerebral trace of declarative memories. *Proc Natl Acad Sci U S A* 2007;104:18778–18783. [PubMed: 18000060]
33. Marshall L, Born J. The contribution of sleep to hippocampus-dependent memory consolidation. *Trends Cogn Sci* 2007;11:442–450. [PubMed: 17905642]
34. Fischer S, Hallschmid M, Elsner AL, Born J. Sleep forms memory for finger skills. *Proc Natl Acad Sci USA* 2002;99:11987–11991. [PubMed: 12193650]
35. Robertson EM, Pascual-Leone A, Press DZ. Awareness modifies the skill-learning benefits of sleep. *Curr Biol* 2004;14:208–212. [PubMed: 14761652]
36. Brown RM, Robertson EM. Off-line processing: reciprocal interactions between declarative and procedural memories. *J Neurosci* 2007;27:10468–10475. [PubMed: 17898218]
37. Krakauer JW, Ghez C, Ghilardi MF. Adaptation to visuomotor transformations: consolidation, interference, and forgetting. *J Neurosci* 2005;25:473–478. [PubMed: 15647491]
38. Huber R, Ghilardi MF, Massimini M, Tononi G. Local sleep and learning. *Nature* 2004;430:78–81. [PubMed: 15184907]
39. Robertson EM, Pascual-Leone A, Miall RC. Current concepts in procedural consolidation. *Nat Rev Neurosci* 2004;5:576–582. [PubMed: 15208699]
40. Goedert KM, Willingham DB. Patterns of interference in sequence learning and prism adaptation inconsistent with the consolidation hypothesis. *Learn Mem* 2002;9:279–292. [PubMed: 12359837]
41. Brashers-Krug T, Shadmehr R, Bizzi E. Consolidation in human motor memory. *Nature* 1996;382:252–255. [PubMed: 8717039]
42. Shadmehr R, Brashers-Krug T. Functional stages in the formation of human long-term motor memory. *J Neurosci* 1997;17:409–419. [PubMed: 8987766]
43. Korman M, Doyon J, Doljansky J, et al. Daytime sleep condenses the time course of motor memory consolidation. *Nat Neurosci* 2007;10:1206–1213. [PubMed: 17694051]
44. Walker MP, Brakefield T, Hobson JA, Stickgold R. Dissociable stages of human memory consolidation and reconsolidation. *Nature* 2003;425:616–620. [PubMed: 14534587]
45. Wright DL, Black CB, Immink MA, et al. Long-term motor programming improvements occur via concatenation of movement sequences during random but not during blocked practice. *J Mot Behav* 2004;36:39–50. [PubMed: 14766487]
46. Immink MA, Wright DL. Motor programming during practice conditions high and low in contextual interference. *J Exp Psychol Hum Percept Perform* 2001;27:423–437. [PubMed: 11318057]
47. Ungerleider LG, Doyon J, Karni A. Imaging brain plasticity during motor skill learning. *Neurobiol Learn Mem* 2002;78:553–564. [PubMed: 12559834]
48. Nezafat R, Shadmehr R, Holcomb HH. Long-term adaptation to dynamics of reaching movements: a PET study. *Exp Brain Res* 2001;140:66–76. [PubMed: 11500799]
49. Grafton ST, Hazeltine E, Ivry RB. Abstract and effector-specific representations of motor sequences identified with PET. *J Neurosci* 1998;18:9420–9428. [PubMed: 9801380]

50. Honda M, Deiber MP, Ibanez V, et al. Dynamic cortical involvement in implicit and explicit motor sequence learning. A PET study. *Brain* 1998;121:2159–2173. [PubMed: 9827775]
51. Classen J, Liepert J, Wise SP, et al. Rapid plasticity of human cortical movement representation induced by practice. *J Neurophysiol* 1998;79:1117–1123. [PubMed: 9463469]
52. Tong C, Flanagan JR. Task-specific internal models for kinematic transformations. *J Neurophysiol* 2003;90:578–585. [PubMed: 12904486]
53. Hadipour-Niktarash A, Lee CK, Desmond JE, Shadmehr R. Impairment of retention but not acquisition of a visuomotor skill through time-dependent disruption of primary motor cortex. *J Neurosci* 2007;27:13413–13419. [PubMed: 18057199]
54. Richardson AG, Overduin SA, Valero-Cabré A, et al. Disruption of primary motor cortex before learning impairs memory of movement dynamics. *J Neurosci* 2006;26:12466–12470. [PubMed: 17135408]
55. Pascual-Leone A, Grafman J, Hallett M. Modulation of cortical motor output maps during development of implicit and explicit knowledge. *Science* 1994;263:1287–1289. [PubMed: 8122113]
56. Robertson EM, Press DZ, Pascual-Leone A. Off-line learning and the primary motor cortex. *J Neurosci* 2005;25:6372–6378. [PubMed: 16000627]
57. Nissen M, Bullemer P. Attentional requirements of learning: evidence from performance measures. *Cognit Psychol* 1987;19:1–32.
58. Plautz EJ, Milliken GW, Nudo RJ. Effects of repetitive motor training on movement representations in adult squirrel monkeys: role of use versus learning. *Neurobiol Learn Mem* 2000;74:27–55. [PubMed: 10873519]
59. Nudo RJ, Milliken GW, Jenkins WM, Merzenich MM. Use-dependent alterations of movement representations in primary motor cortex of adult squirrel monkeys. *J Neurosci* 1996;16:785–807. [PubMed: 8551360]
60. Duque J, Murase N, Celnik P, et al. Intermanual Differences in movement-related interhemispheric inhibition. *J Cogn Neurosci* 2007;19:204–213. [PubMed: 17280510]
61. Reis J, Swayne OB, Vandermeeren Y, et al. Contribution of transcranial magnetic stimulation to the understanding of cortical mechanisms involved in motor control. *J Physiol* 2008;586:325–351. [PubMed: 17974592]
62. Ward NS, Cohen LG. Mechanisms underlying recovery of motor function after stroke. *Arch Neurol* 2004;61:1844–1848. [PubMed: 15596603]
63. Avanzino L, Teo JT, Rothwell JC. Intracortical circuits modulate transcallosal inhibition in humans. *J Physiol* 2007;583:99–114. [PubMed: 17556392]
64. Perez MA, Cohen LG. Mechanisms underlying functional changes in the primary motor cortex ipsilateral to an active hand. *J Neurosci*. 2008 (in press).
65. Kobayashi M, Hutchinson S, Théoret H, et al. Repetitive TMS of the motor cortex improves ipsilateral sequential simple finger movements. *Neurology* 2004;62:91–98. [PubMed: 14718704]
66. Schambra HM, Sawaki L, Cohen LG. Modulation of excitability of human motor cortex (M1) by 1 Hz transcranial magnetic stimulation of the contralateral M1. *Clin Neurophysiol* 2003;114:130–133. [PubMed: 12495773]
67. Plewnia C, Lotze M, Gerloff C. Disinhibition of the contralateral motor cortex by low-frequency rTMS. *Neuroreport* 2003;14:609–612. [PubMed: 12657896]
68. Wassermann EM, Wedgegartner FR, Ziemann U, et al. Crossed reduction of human motor cortex excitability by 1-Hz transcranial magnetic stimulation. *Neurosci Lett* 1998;250:141–144. [PubMed: 9708852]
69. Daskalakis ZJ, Paradiso GO, Christensen BK, et al. Exploring the connectivity between the cerebellum and motor cortex in humans. *J Physiol* 2004;557:689–700. [PubMed: 15047772]
70. Koch G, Franca M, Mochizuki H, et al. Interactions between pairs of transcranial magnetic stimuli over the human left dorsal premotor cortex differ from those seen in primary motor cortex. *J Physiol* 2007;578:551–562. [PubMed: 17124263]
71. Nitsche MA, Paulus W. Sustained excitability elevations induced by transcranial DC motor cortex stimulation in humans. *Neurology* 2001;57:1899–1901. [PubMed: 11723286]

72. Kim YH, Park JW, Ko MH, et al. Facilitative effect of high frequency subthreshold repetitive transcranial magnetic stimulation on complex sequential motor learning in humans. *Neurosci Lett* 2004;367:181–185. [PubMed: 15331148]
73. Vines BW, Nair DG, Schlaug G. Contralateral and ipsilateral motor effects after transcranial direct current stimulation. *Neuroreport* 2006;17:671–674. [PubMed: 16603933]
74. Antal A, Nitsche M, Kincses T, et al. Facilitation of visuo-motor learning by transcranial direct current stimulation of the motor and extrastriate visual areas in humans. *Eur J Neurosci* 2004;19:2888–2892. [PubMed: 15147322]
75. Nitsche M, Schauenburg A, Lang N, et al. Facilitation of implicit motor learning by weak transcranial direct current stimulation of the primary motor cortex in the human. *J Cogn Neurosci* 2003;15:619–626. [PubMed: 12803972]
76. Jebsen RH, Taylor N, Trieschmann RB, et al. An objective and standardized test of hand function. *Arch Phys Med Rehabil* 1969;50:311–319. [PubMed: 5788487]
77. Boggio P, Castro L, Savagim E, et al. Enhancement of non-dominant hand motor function by anodal transcranial direct current stimulation. *Neurosci Lett* 2006;404:232–236. [PubMed: 16808997]
78. Gilio F, Rizzo V, Siebner HR, Rothwell JC. Effects on the right motor hand-area excitability produced by low-frequency rTMS over human contralateral homologous cortex. *J Physiol* 2003;551:563–573. [PubMed: 12821724]
79. Agostino R, Iezzi E, Dinapoli L, et al. Effects of 5 Hz subthreshold magnetic stimulation of primary motor cortex on fast finger movements in normal subjects. *Exp Brain Res* 2007;180:105–111. [PubMed: 17265042]
80. Carey JR, Fregni F, Pascual-Leone A. rTMS combined with motor learning training in healthy subjects. *Restor Neurol Neurosci* 2006;24:191–199. [PubMed: 16873974]
81. Siebner HR, Lang N, Rizzo V, et al. Preconditioning of low-frequency repetitive transcranial magnetic stimulation with transcranial direct current stimulation: evidence for homeostatic plasticity in the human motor cortex. *J Neurosci* 2004;24:3379–3385. [PubMed: 15056717]
82. Silvanto J, Muggleton NG, Cowey A, Walsh V. Neural activation state determines behavioral susceptibility to modified theta burst transcranial magnetic stimulation. *Eur J Neurosci* 2007;26:523–528. [PubMed: 17650122]
83. Ziemann USHR. Modifying motor learning through gating and homeostatic metaplasticity. *Brain Stimulation* 2008;1:60–66.
84. Ziemann U, Ilic TV, Jung P. Long-term potentiation (LTP)-like plasticity and learning in human motor cortex--investigations with transcranial magnetic stimulation (TMS). *Suppl Clin Neurophysiol* 2006;59:19–25. [PubMed: 16893088]
85. Gentner R, Wanklerl K, Reinsberger C, et al. Depression of Human Corticospinal Excitability Induced by Magnetic Theta-burst Stimulation: Evidence of Rapid Polarity-Reversing Metaplasticity. *Cereb Cortex*. 2007