

# Targeting brain networks with multichannel transcranial current stimulation (tCS)

Giulio Ruffini<sup>1,2</sup>, Fabrice Wendling<sup>3</sup>, Roser Sanchez-Todo<sup>1</sup> and Emiliano Santarnecchi<sup>4</sup>

## Abstract

The brain is a complex, plastic, electrical network whose dysfunctions result in neurological disorders. Multichannel transcranial electrical stimulation (tCS) is a non-invasive neuromodulatory technique with the potential for network-oriented therapy. Challenges to realizing this vision include the proper identification of involved networks in a patient-specific context, a deeper understanding of the effects of stimulation on interconnected neuronal populations - both immediate and plastic - and, based on these, developing strategies to personalize brain stimulation interventions. For this reason, personalized *hybrid* biophysical and physiological models of brain networks are poised to play a key role in the evolution of network-oriented transcranial stimulation. We review some of the recent work in this emerging area of research and provide an outlook for future modeling and experimental work, as well as for developing its clinical applications in fields such as epilepsy.

## Addresses

<sup>1</sup> Neuroelectronics, 08035 Barcelona, Spain

<sup>2</sup> Starlab Barcelona, 08035 Barcelona, Spain

<sup>3</sup> LTSI, Univ-Rennes, INSERM, Rennes, France

<sup>4</sup> Berenson-Allen Center for Non-Invasive Brain Stimulation, Beth Israel Medical Center, Harvard Medical School, Boston, MA, USA

Corresponding author: Ruffini, Giulio ([giulio.ruffini@neuroelectronics.com](mailto:giulio.ruffini@neuroelectronics.com))

Current Opinion in Biomedical Engineering 2018, 8:70–77

This review comes from a themed issue on **Neural Engineering: Neuromodulation**

Edited by Aysegul Gunduz, Giulio Ruffini, Christine Schmidt and Jose De Millan

Received 9 October 2018, revised 13 October 2018, accepted 13 November 2018

<https://doi.org/10.1016/j.cobme.2018.11.001>

2468-4511/© 2018 Elsevier Inc. All rights reserved.

## Keywords

tDCS, tCS, tACS, tES, fMRI, rs-fcMRI.

## Introduction

The brain is a complex, plastic, electrical network operating at multiple scales - neural processing is essentially mediated by functional and structural networks. Over the past decades, neuroscience has made significant advances in our understanding of brain function. There is a

growing body of evidence suggesting that large-scale networks underlie both integration and differentiation processes which are fundamental for information processing in the brain. For instance, putatively simple cognitive tasks such as object recognition have been shown to involve networks that include the bilateral occipital, the left temporal and the left/right frontal regions [1]. Neuropsychiatric disorders ultimately result from network dysfunctions which may arise from the abnormality in one or more isolated brain regions but produce alterations in larger brain networks (see Refs. [2–4] and references therein).

In such a context, networks become the natural target of neuromodulatory interventions. Advances in neuroimaging modalities such as positron emission tomography (PET), magneto- and electroencephalography (EEG/MEG), functional magnetic resonance imaging (fMRI) and diffusion tensor imaging (DTI) provide valuable tools for the identification of networks. For instance, the ‘resting state’ paradigm is increasingly used to assess intrinsic brain activity and brain connectivity using modalities such as fMRI, EEG or MEG [5]. Activity recorded during spontaneous rest using fMRI can be decomposed into separate but integrated resting-state networks (RSNs) [6,7] also known as “modules” [8] or “architectures” [9], with specific RSNs reflecting the activity within sensory (e.g., visual, motor, auditory) and associative brain regions related to high-order cognitive processes such as abstract reasoning, attention, language, and memory. This organization, as captured via functional connectivity (FC) analysis of fMRI data collected during resting-state (rs-fcMRI), is correlated with individual variability in several cognitive functions and personality traits [10–13], with recent studies suggesting the possibility of capturing individual brain uniqueness by means of finely tailored FC analysis [14]. A similar approach can be used for the spatio-temporal decomposition of electrophysiological signals at higher temporal resolution (~1 ms) into so-called microstates [15,16]. These have been linked to a variety of cognitive functions and pathologies [16–19]. While such approaches provide stimulation targets at relatively high spatial resolution, currently used noninvasive brain stimulation techniques - such as Transcranial Magnetic Stimulation (TMS) - cannot easily be employed to simultaneously engage multiple network nodes or sub-networks. TMS network manipulation based on the induction of spike-timing-dependent

plasticity (STDP) has been recently proposed [20,21], but requires relatively expensive hardware and can only be performed in laboratory settings. Novel, safe, portable solutions for network-engagement are needed.

Transcranial electrical current stimulation (*tCS*, sometimes also called *tES*), which includes both direct and alternating current variants known as tDCS and tACS, is a non-invasive sub-threshold neuromodulatory technique pioneered by Nitsche and Paulus [22]. Low intensity, controlled currents (typically  $\sim 1$  mA but  $< 4$  mA) are applied through scalp electrodes in repeated 20–40-min sessions. This subtle but persistent modulation of neuronal activity is believed to lead to plastic effects deriving from Hebbian mechanisms (see Refs. [23–25] and references therein). That is, tCS induces concurrent and plastic effects from persistent (in time), mesoscale (in space), weak electric fields acting on brain networks. Its clinical applications include neuropathic chronic pain, major depression, stroke rehabilitation, addictive disorders and epilepsy [26]. tCS is recognized for its applicability and safety [27].

The recent evolution of tCS has delivered multichannel systems using small electrodes much like EEG. This advance comes with opportunities and challenges.

### Multichannel stimulation and its optimization

In the past years, methods have been proposed to optimize multichannel tCS (see, e.g. Ref. [28]) proved that this problem is mathematically well-posed and showed that optimized electric fields display significantly higher focality and, in general, a better alignment with the target vector than those produced by standard bipolar electrode montages [29]. Based on work from Miranda et al. [30] and Fox et al. [31], Ruffini et al. [32] proposed a method for optimization of multichannel tCS (*Stimweaver*). Its main features are a focus on cortical excitability, the use of an interaction mechanism inferred from prior *in-vivo* and *in-vitro* work (called the *lambda-E model* [23]) that places emphasis on the component of the electric field orthogonal to the cortical surface, MRI driven finite element modeling of the electric fields produced by multichannel tCS, and a rapid optimization method exploring number, current intensity and spatial location of electrodes. The method requires as key inputs a specification of the target electric field on the cortex, a weight map to prioritize target regions for the optimizer and other parameters such as the maximal number of electrodes and currents allowed. Defining these maps is, of course, key and requires a deep understanding of cortical function – including its network aspects.

This method has been employed by several research groups. For example, Fisher et al. [33] explored whether

the effects of tCS on a region can be enhanced by targeting its associated network. In particular, a network associated with a local target on the left motor cortex (M1) was defined using rs-fcMRI. In a cross-over study, fifteen healthy subjects were stimulated in several conditions, including one with a bipolar montage targeting the seed (M1), another with an eight-electrode montage targeting its associated resting state network, and a sham condition. Cortical excitability of the left M1 was probed using TMS/MEPs, as in the pioneering work by Nitsche and Paulus [22]. The authors observed that network-targeted tDCS led to a significant increase in left M1 excitability over time compared to traditional tDCS.


Dagan et al. [34] recently studied the use of multichannel tDCS in Parkinson's disease (PD) with freezing of gait (FOG), one of its most disturbing and least understood symptoms. Several hypotheses suggest that FOG is not only a motor problem but also partly the result of deficits in executive function mediated by the dorsolateral prefrontal cortex (DLPFC) (see Ref. [34] and references therein). Indeed, targeting the DLPFC with tDCS appears to positively affect cognition, gait, and postural control in other populations. Because PD manifests strongly as a motor disturbance phenomenon, including FOG, most studies in PD have also focused on M1, reporting motor function and gait improvements with bipolar tDCS compared to sham stimulation (see references in Ref. [34]). Dagan et al. [34] employed multichannel tCS optimized for maximizing facilitation of both primary motor cortex (M1) and the left dorsolateral prefrontal cortex (DLPFC), and compared this to stimulation of M1 only and a Sham condition. Multi-target stimulation of both areas provided a significant improvement over the other conditions.

In another recent example, research in disorders of consciousness has employed multichannel network stimulation [35] sought to engage the external (frontoparietal) consciousness network in severely brain-injured patients using a target map derived from rs-fMRI. Finally, in yet another example with healthy subjects, attempts have been made to optimize multichannel solutions engaging cortical networks relevant for cognitive training, e.g., targeting flexibility or working memory-related nodes while participants were undergoing executive function training [36].

### tCS and brain networks: from biophysics to physiology

If the critical features of pathological networks can be effectively captured in computational models, they can be used for diagnosis and delivery of personalized therapeutic weak electric fields (Figure 1). Multiple studies in theoretical and computational neuroscience

Figure 1




**Workflow for the creation of a biophysical model and for model-driven tCS optimization.** Anatomical MRI data (1) is used to create a finite element biophysical model (FEM), and electrodes are placed using the 10-10 EEG system (2) (see Miranda et al., 2018 [37] for a review). A target specification is provided, together with the desired number of electrodes and maximal currents. The *Stimweaver* algorithm provides the solution (3, i.e., electrode positions and currents). The approach is applicable to tDCS, tACS and other tCS modalities [32].

have developed whole-brain network models [38–41] to explore the relationship between brain function and its underlying connectivity. This increased interest in finding the origin of the structure-function relationship has led to a newly developing field known as *network neuroscience* (Bassett and Sporns, 2017) [42] that relies on graph theory to study the brain across its multiple scales and complexities. Following earlier work by Merlet et al. [39], Sanchez-Todo et al. [43] develop a method that allows for the use of a subject's EEG and MRI for the creation of a personalized whole brain model. The model is optimized to reproduce a subject's EEG and allows for virtual brain stimulation, and hence optimization (see Figure 2). Earlier, Bansal et al. [44], Spiegler et al. [45], and Muldoon et al. [46] proposed a similar approach. Although “hybrid” models can produce physiologically-plausible EEG and simulate the generation of realistic tCS electric fields (see Miranda et al. [37] in this issue, and references therein), representing faithfully the effects of neuromodulation on brain plasticity remains an unresolved, important challenge. We now know from experimental work that tCS can directly

impact neuronal excitability and synaptic plasticity [47,48]. Marquez-Ruiz et al. [49], e.g., showed that blocking adenosine A1 receptors prevents the long-term depression evoked in the somatosensory cortex after cathodal tDCS in the rabbit. Based on molecular and functional investigations (immunoblotting, immunofluorescence, and electrophysiological recordings), Paciello et al. [50] provide novel evidence that anodal tDCS affects structural plasticity of the rat auditory cortex in a paradigm of noise-induced hearing loss. Wischniewski et al. [51] also reported that 20 Hz tACS can alter NMDA Receptor-Mediated plasticity in the human motor cortex. A number of studies indicate that tCS can alter the release of neurotransmitters, typically dopamine [52], glutamate and GABA [53]. Ultimately, electric field mediated effects translate into short- or long-term changes in the network connectivity - and therapeutic effects. As the precise mechanisms involved in tCS-induced plasticity changes still remain elusive, multiscale computational models offer a unique framework to untangle them, allowing, for instance, to distinguish effects occurring at presynaptic (membrane

Figure 2



**Workflow for the creation of hybrid models model-driven tCS optimization.** DTI and anatomical MRI data are combined to create a finite element biophysical model (FEM), which is then personalized using EEG and other data to reflect *both* biophysical and physiologic characteristics – from excitation/inhibition balance to plastic potential (long-term effects physiological model). The personalized hybrid brain model can be used to generate EEG and to simulate the effects of brain stimulation. As a result, personalized diagnosis and treatment are possible, including the optimization of stimulation protocols.

polarization of axon terminals, neurotransmitter release) or postsynaptic (GABA or glutamate receptors) level.

## Future applications

### Epilepsy

Epilepsy is a devastating, chronic disease that severely affects the quality of life of 65 million people worldwide (WHO, Fact Sheet on Epilepsy, 2012), 35% of whom do not respond to drugs. Almost a third of patients (29%) are untreatable: in about 19 million patients, drugs fail, and surgery is not an option or has failed too. Treatment-resistant epilepsies represent not only a considerable challenge for the health care system but also a tremendous burden at the individual, family, and community levels [54]. They are characterized by an epileptogenic network (EN) interconnecting distant brain areas located in one of the two hemispheres. There is a large body of evidence suggesting that patient-specific ENs [55] are responsible for the generation and spread of seizures through synchronization processes that interconnect neuronal assemblies with altered excitability [56]. Of note, some studies have tried to predict surgical outcome by removing EN edges of the patient specific connectivity data in computational models of the subject's brain [57–61]. In this context, tCS can represent a valuable alternative to surgery [62], provided that fundamental issues are addressed. First, epileptogenic networks are patient-specific. Therefore, interventions must be “tailored” to each patient based on the accurate definition of target brain areas and networks. Second, stimulation protocols must achieve a therapeutic effect through a “network-


aware” management of hyperexcitability - a hallmark of epileptogenic systems. Third, therapeutic effects must be optimized in order to prevent the occurrence of seizures. A protective and durable effect will certainly require a better understanding of the mechanisms of action of weak electric fields on brain networks from short (minutes to hours) to long (days, weeks) time scales.

### Reaching subcortical targets via networks

Fox et al. [63] identified diseases treated with both non-invasive and deep brain stimulation (DBS), listed the target sites thought to be most effective in each disease and tested the hypothesis that these sites are nodes within a brain network as defined by rs-fcMRI. They found that sites in which DBS was effective were functionally connected to sites where noninvasive brain stimulation had been found to be effective in diseases including depression, PD, obsessive-compulsive disorder, essential tremor, addiction, pain, minimally conscious state, and Alzheimer's disease. This suggests that rs-fcMRI may be useful for translating therapy across stimulation modalities, optimizing treatment, and for the identification of new stimulation targets. It also supports a more general network approach toward understanding and treating neuropsychiatric disease, highlighting the therapeutic potential of targeted brain network modulation. Examples of potential cortical and subcortical targets relevant for neuropsychiatric conditions, as well as their rs-fcMRI map and corresponding multichannel optimization, are shown in Figure 3 (see also Ruffini et al. [32] for further discussion on the use of these maps for multichannel tCS optimization).



Figure 3



**Connectivity-based network targeting.** (A) Cortical representation of rs-fMRI connectivity patterns of selected brain regions of clinical relevance in various neuropsychiatric conditions. The target regions are used as seeds, and their pattern of positively and negatively correlated regions in the brain are computed. Multichannel tCS can be optimized to enhance positive (“excitatory”) or negative (“inhibitory”) cortical nodes, inducing changes in their connectivity with the seed region and possibly modulating its spontaneous activity. (B) Examples of multichannel tCS solutions derived from *Stimweaver* [32] for targeting the supplementary motor area in patients with obsessive-compulsive disorder (OCD) and subgenual cortex in patients with depression.

## Stroke

In another example, Otal et al. [64] proposed to identify networks affected by a stroke at the individual level: location, extent, and pattern of functional network connectivity disruption should be considered when determining the optimal tDCS intervention. Alstott et al. [65] did a related *in silico* study where network edges were removed to investigate the effect of such perturbations on simulated brain activity. See an extended review of Aerts et al. [66] regarding computational lesion and empirical studies investigating brain network alterations in cancer, stroke and traumatic

injury patients. In the case of stroke, each lesion type displays a particular functional and structural connectivity signature that determines the tDCS intervention goals. Lesion topography is usually subcortical, with intracortical connectivity disruptions contributing strongly to behavioral deficits [67]. In general, we may consider three main approaches: a) targeting a single region or node, b) targeting the single region indirectly via a network as described above, or c) select a network or sub-network (i.e., multi-nodal) as the target. The latter may be especially relevant given the correlation of connectivity disruption and symptoms. Depending on the approach

chosen, different optimization strategies can be envisioned. Tools based on tractography can be used to assess damaged networks and devise therapeutic strategies [68].

## Conclusions

Multichannel tCS provides a promising tool for targeting networks but is not yet used as a standard treatment in any disease. This is related to several challenges and methodological limitations: an overwhelming number of stimulation parameter combinations, empirical parameter setting, an absence of a rational definition of targets and protocols, the qualitative nature of results, unknown mechanisms of action, and an insufficient account for patient-specific factors. A bottom-up, science-based mechanistic understanding of both the effects of tCS and the desired cortical network changes is lacking. Research should aim to overcome this by providing a better understanding mechanism of interaction - including both immediate and longer-term plastic effects of electric fields in networks across scales. We also need to refine the current methods used for network identification and for the design of interventions in each disease and patient. Finally, experiments should be carried out to investigate how network interactions can best be leveraged by tCS, measuring the functional and structural alterations induced by tCS using tools such as EEG, fMRI or DTI. Modeling in sufficient detail the combined biophysics and physiology of tCS will be paramount for the design and interpretation of studies and for their ultimate clinical translation.

## Acknowledgments

This research was supported in part by the Future Emerging Technologies Open Luminous project (H2020-FETOPEN-2014-2015-RIA under agreement No. 686764) as part of the European Union's Horizon 2020 research and training program 2014–2018. ES is partially supported by Office of the Director of National Intelligence (ODNI), Intelligence Advanced Research Projects Activity, via 2014-13121700007. The views and conclusions contained herein are those of the authors and should not be interpreted as necessarily representing the official policies or endorsements, either expressed or implied, of the ODNI, IARPA, or the U.S. Government. ES is supported by the Beth Israel Deaconess Medical Center via the Chief Academic Officer (CAO) Award 2017, and the Defense Advanced Research Projects Agency (DARPA) via HR001117S0030. The content of this paper is solely the responsibility of the authors and does not necessarily represent the official views of Harvard University, and its affiliated academic health care centers.

## Conflict of interest statement

Giulio Ruffini is a shareholder and works for Neuro-electrics, a company designing medical devices for brain stimulation. Roser Sanchez-Todo is a researcher at Neuroelectronics.

## References

Papers of particular interest, published within the period of review, have been highlighted as:

- \* of special interest
- \*\* of outstanding interest

1. Price C, Moore C, Humphreys G, Frackowiak R, Friston K: **The neural regions sustaining object recognition and naming.** *Proc R Soc Lond Ser B: Biol Sci* 1996, **263**:1501–1507.
2. Fox Michael D, *et al.*: **Measuring and manipulating brain connectivity with resting state functional connectivity magnetic resonance imaging (fcMRI) and transcranial magnetic stimulation (TMS).** *Neuroimage* 2012a, **62**:4.
3. Fox MD, Buckner RL, White MP, Greicius MD, Pascual-Leone A: **Efficacy of transcranial magnetic stimulation targets for depression is related to intrinsic functional connectivity with the subgenual cingulate.** *Biol Psychiatry* 2012b, **72**.
4. Fornito A, Zalesky A, Breakspear M: **The connectomics of brain disorders.** *Nat Rev Neurosci* 2015, **16**:159–172. <https://doi.org/10.1038/nrn3901>.
5. Van Diessen E, Numan T, van Dellen E, van der Kooij A, Boersma M, Hofman D, *et al.*: **Opportunities and methodological challenges in EEG and MEG resting state functional network research.** *Clin Neurophysiol* 2015, **126**:1468–1481.
6. Achard S, Bullmore E: **Efficiency and cost of economical brain functional networks.** *PLoS Comput Biol* 2007, **3**:e17.
7. Sporns O: **The non-random brain: efficiency, economy, and complex dynamics.** *Front Comput Neurosci* 2011, **5**:5. <https://doi.org/10.3389/fncom.2011.00005>.
8. Power JD, Cohen AL, Nelson SM, Wig GS, Barnes KA, Church JA, Vogel AC, Laumann TO, Miezin FM, Schlaggar BL, Petersen SE: **Functional network organization of the human brain.** *Neuron* 2011, **72**:665–678. <https://doi.org/10.1016/j.neuron.2011.09.006>.
9. Hearne LJ, Cocchi L, Zalesky A, Mattingley JB: **Reconfiguration of brain network architectures between resting state and complexity-dependent cognitive reasoning.** *J Neurosci* 30 August 2017, **37**:8399–8411. <https://doi.org/10.1523/JNEUROSCI.0485-17.2017>.
10. Adelstein JS, Shehzad Z, Mennes M, Deyoung CG, Zuo XN, Kelly C, Margulies DS, Bloomfield A, Gray JR, Castellanos FX, Milham MP: **Personality is reflected in the brain's intrinsic functional architecture.** *PLoS One* 2011, **6**:e27633.
11. Corbetta M, Shulman GL: **Control of goal-directed and stimulus-driven attention in the brain.** *Nat Rev Neurosci* 2002, **3**:201–215. <https://doi.org/10.1038/nrn755>.
12. Santarnecchi E, Emmendorfer A, Tadayon S, Rossi S, Rossi A, Pascual-Leone A: **Network connectivity correlates of variability in fluid intelligence performance.** *Intelligence* 2017a, **65**: 35–47. <https://doi.org/10.1016/j.intell.2017.10.002>.
13. Santarnecchi E, Sprugnoli G, Tatti E, Mencarelli L, Neri F, Momi D, Di Lorenzo G, Pascual-Leone A, Rossi S, Rossi A: **Brain functional connectivity correlates of coping styles.** *Cognit Affect Behav Neurosci* 2018. <https://doi.org/10.3758/s13415-018-0583-7>.
14. Finn ES, Shen X, Scheinost D, Rosenberg MD, Huang J, Chun MM, Papademetris X, Constable RT: **Functional connectome fingerprinting: identifying individuals using patterns of brain connectivity.** *Nat Neurosci* 2015, **18**:1664–1671. <https://doi.org/10.1038/nn.4135>.
15. Khanna A, Pascual-Leone A, Michel CM, Farzan F: **Microstates in resting-state EEG: current status and future directions.** *Neurosci Biobehav Rev* 2015, **49**:105–113. <https://doi.org/10.1016/j.neubiorev.2014.12.010>.
16. Lehmann D, Strik WK, Henggeler B, Koenig T, Koukkou M: **Brain electric microstates and momentary conscious mind states as building blocks of spontaneous thinking: I. Visual imagery and abstract thoughts.** *Int J Psychophysiol Off J Int Organ Psychophysiol* 1998, **29**:1–11.
17. Andreou C, Faber PL, Leicht G, Schoettle D, Polomac N, Hanganu-Opatz IL, Lehmann D, Mulert C: **Resting-state connectivity in the prodromal phase of schizophrenia: insights from EEG microstates.** *Schizophr Res* 2014, **152**:513–520. <https://doi.org/10.1016/j.schres.2013.12.008>.
18. Nishida K, Morishima Y, Yoshimura M, Isotani T, Irisawa S, Jann K, Dierks T, Strik W, Kinoshita T, Koenig T: **EEG microstates associated with salience and frontoparietal networks in frontotemporal dementia, schizophrenia and Alzheimer's disease.** *Clin Neurophysiol* 2013, **124**:1106–1114. <https://doi.org/10.1016/j.clinph.2013.01.005>.

19. Santarnecchi E, Khanna AR, Musaeus CS, Benwell CSY, Davila P, Farzan F, Matham S, Pascual-Leone A, Shafi MM, On behalf of Honeywell SHARP Team authors: **EEG microstate correlates of fluid intelligence and response to cognitive training.** *Brain Topogr* 2017. <https://doi.org/10.1007/s10548-017-0565-z>.
20. Santarnecchi E, Momi D, Sprugnoli G, Neri F, Pascual-Leone A, Rossi A, Rossi S: **Modulation of network-to-network connectivity via spike-timing-dependent noninvasive brain stimulation.** *Hum Brain Mapp* 2018. <https://doi.org/10.1002/hbm.24329>.
21. Veniero D, Ponzo V, Koch G: **Paired associative stimulation enforces the communication between interconnected areas.** *J Neurosci* 2013, **33**:13773–13783. <https://doi.org/10.1523/JNEUROSCI.1777-13.2013>.
22. Nitsche MA, Paulus W: **Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation.** *J Physiol* 2000, **527**:633–639.
23. Ruffini G, Wendling F, Merlet I, Molae-Ardekani B, Mekonnen A, Salvador R, Soria-Frisch A, Grau C, Dunne S, Miranda PC: **Transcranial current brain stimulation (tCS): models and technologies.** *IEEE Trans Neural Syst Rehabil Eng* 2013, **21**: 333–345.
24. Henrich-noack P, Sergeeva EG, Sabel BA: **Non-invasive electrical brain stimulation: from acute to late-stage treatment of central nervous system damage.** *Neural Regen Res* 2017, **12**. <https://doi.org/10.4103/1673-5374.217322Nitsche>.
25. Karabanov A, Ziemann U, Hamada M, George MS, Quartarone A, Classen J, et al.: **Brain stimulation consensus paper: probing homeostatic plasticity of human cortex with non-invasive transcranial brain stimulation.** *Brain Stimul* 2015, **8**:442–454. <https://doi.org/10.1016/j.brs.2015.01.404>.
26. Lefaucheur JP, Antal A, Ayache SS, Benninger DH, Brunelin J, Cogiamanian F, Cotelli M, De Ridder D, Ferrucci R, Langguth B, Marangolo P, Mylius V, Nitsche MA, Padberg F, Palm U, Poulet E, Priori A, Rossi S, Schecklmann M, Vanneste S, Ziemann U, Garcia-Larrea L, Paulus: **Evidence-based guidelines on the therapeutic use of transcranial direct current stimulation (tDCS).** *Clin Neurophysiol* 2017, **128**:56–92. <https://doi.org/10.1016/j.clinph.2016.10.087>. Epub 2016 Oct 29.
- This paper provides an up to date review of the clinical uses of tDCS, highlighting those applications that show probable efficacy.
27. Antal A, Alekseichuk I, Bikson M, Brockmüller J, Brunoni AR, Chen R, Cohen LG, Dowthwaite G, Ellrich J, Flöel A, Fregni F, George MS, Hamilton R, Hauelsen J, Herrmann CS, Hummel FC, Lefaucheur JP, Liebetanz D, Loo CK, McCaig CD, Miniussi C, Miranda PC, Moliadze V, Nitsche MA, Nowak R, Padberg F, Pascual-Leone A, Poppendieck W, Priori A, Rossi S, Rossini PM, Rothwell J, Rueger MA, Ruffini G, Schellhorn K, Siebner HR, Ugawa Y, Wexler A, Ziemann U, Hallett M, Paulus W: **Low intensity transcranial electric stimulation: safety, ethical, legal regulatory and application guidelines.** *Clin Neurophysiol* 2017, **128**:1774–1809. <https://doi.org/10.1016/j.clinph.2017.06.001>.
28. Dmochowski JP, Datta A, Bikson M, Su Y, Parra LC: **Optimized multi-electrode stimulation increases focality and intensity at target.** *J Neural Eng* 2011, **8**:046011. <https://doi.org/10.1088/1741-2560/8/4/046011>.
29. Wagner S, Burger M, Wolters CH: **An optimization approach for well-targeted transcranial direct current stimulation.** *SIAM J Appl Math* 2016, **76**:2154–2174. <https://doi.org/10.1137/15m1026481>.
30. Miranda PC, Mekonnen A, Salvador R, Ruffini G: **The electric field in the cortex during transcranial current stimulation.** *Neuroimage* 2013, **70**:48–58.
31. Fox Michael D, Buckner Randy L, Liu Hesheng, Chakravarty MM, Lozano Andres M, Pascual-Leone Alvaro: **Resting-state networks link invasive and noninvasive brain stimulation across diverse psychiatric and neurological diseases.** *Proc Natl Acad Sci USA* 2014b, **111**:E4367–E4375.
32. Ruffini G, Fox MD, Ripolles O, Miranda PC, Pascual-Leone A: **Optimization of multifocal transcranial current stimulation for weighted cortical pattern targeting from realistic modeling of electric fields.** *Neuroimage* 2014, **89**:216–225.
33. Fisher DB, et al.: **Network-targeted non-invasive brain stimulation with multifocal tDCS.** *Brain Stimul: Basic Transl Clin Res Neuromodulation* 2017, **10**:2.
- This paper evaluates for the first time the potential of seed-based (rs-fMRI) targeting and network stimulation using multichannel tCS. The authors find an enhanced effect of multichannel network vs. bipolar tDCS on motor cortex cortical excitability as measured by TMS/MEPs.
34. Dagan M, Herman T, Harrison R, et al.: **Multitarget transcranial direct current stimulation for freezing of gait in Parkinson's disease.** *Mov Disord: Off J Mov Disord Soc* 2018, **33**:642–646. <https://doi.org/10.1002/mds.27300>.
- This paper proposes a dual target approach to target networks relevant to freezing of gait and related outcomes. By targeting nodes involved in motor and cognitive tasks, the authors show it is possible to improve potential cognitively coupled motor tasks in freezing of gait. The authors also demonstrate the potential of an "active sham" concept using multichannel stimulation for better blinding.
35. Thibaut A, Martens G, Laureys S: **Multichannel tDCS of the frontoparietal network in patients with disorders of consciousness: a double blind sham controlled randomized clinical trial.** *Brain Stimul: Basic Transl Clin Res Neuromodulation* 2017, **10**.
36. Brem A-K, Almquist JN-F, Mansfield K, Plessow F, Sella F, Santarnecchi E, Orhan U, McKanna J, Pavel M, Mathan S, Yeung N, Pascual-Leone A, Kadosh RC, Honeywell SHARP Team authors: **Modulating fluid intelligence performance through combined cognitive training and brain stimulation.** *Neuropsychologia* 2018. <https://doi.org/10.1016/j.neuropsychologia.2018.04.008>.
37. Miranda PC, Callejón-Leblic MA, Salvador R, Ruffini G: **Realistic modeling of transcranial current stimulation: the electric field in the brain.** *Curr Opin Biomed Eng* 2018. <https://doi.org/10.1016/j.cobme.2018.09.002>.
- A review of the state of the art in the modeling of electric fields in the brain, techniques, and validation.
38. Deco G, Jirsa VK, McIntosh AR: **Emerging concepts for the dynamical organization of resting-state activity in the brain.** *Nat Rev Neurosci* 2011, **12**:43–56.
39. Merlet I, Birot G, Salvador R, Molae-Ardekani B, Mekonnen A, Soria-Frisch A, Ruffini G, Miranda PC, Wendling F: **From oscillatory transcranial current stimulation to scalp EEG changes: a biophysical and physiological modeling study.** *PLoS One* 2013, **8**:e57330. <https://doi.org/10.1371/journal.pone.0057330>.
40. Cabral J, Luchkoo H, Woolrich M, Joensson M, Mohseni H, Baker A, ... Deco G: **Exploring mechanisms of spontaneous functional connectivity in MEG: how delayed network interactions lead to structured amplitude envelopes of band-pass filtered oscillations.** *Neuroimage* 2014, **90**:423–435.
41. Bansal K, Johan Nakuci, Sarah Feldt Muldoon: **Personalized brain network models for assessing structure–function relationships.** *Curr Opin Neurobiol* 2018a, **52**:42–47.
- This paper proposes the combination of structural data with neural mass models to study structure-function relationships in the human brain, with applications for therapy (surgery or brain stimulation).
42. Bassett DS, Sporns O: **Network neuroscience.** *Nat Neurosci* 2017, **20**:353–364.
- The authors review emerging trends in network neuroscience and provide a path toward a better understanding of the brain as a multi-scale networked system, and on how to leverage it.
43. Sanchez-Todo R, Salvador R, Santarnecchi E, Wendling F, Deco G, Ruffini G: **Personalization of hybrid brain models from neuroimaging and electrophysiology data.** *BioRxiv* 2018. <https://doi.org/10.1101/461350>.
- This methods paper shows, using real MRI and EEG data, how to integrate structural, anatomical and physiological data to create realistic brain models of the human brain, with applications in brain stimulation and basic neuroscience. Such hybrid models of biophysics and physiology can be used to model optimize tDCS on brain networks, for example.
44. Bansal K, Medaglia J, Bassett D, Vettel J, Muldoon S: **Data-driven brain network models differentiate variability across language tasks.** *PLoS Comput Biol* 2018b, **14**, e1006487. 018.
45. Spiegler A, Hansen ECA, Bernard C, McIntosh AR, Jirsa VK: **Selective activation of resting state networks following focal**



stimulation in a connectome-based network model of the human brain. *eNeuro* 2016, **3**:1–17.

46. Muldoon SF, Pasqualetti F, Gu S, Cieslak M, Grafton ST, Vettel JM, Bassett DS: **Stimulation based control of dynamic brain networks**. *PLoS Comput Biol* 2016, **12**, e1005076.

The authors performed *in silico* experiments in data-driven brain network models to relate patterns of activation due to targeted stimulation. Furthermore, they use their model to explore the relationship between cognitive systems and the underlying brain anatomy.

47. Modolo J, Denoyer Y, Fabrice W, Benquet P: **Physiological effects of low-magnitude electric fields on brain activity: advances from in vitro, in vivo and in silico models**. *Curr Opin Biomed Eng* 2018.

An up to date review on models for the interaction of weak electric fields with neurons and their larger scale effects.

48. Sánchez-León CA, Sánchez-López A, Ammann C, Cordones I, Carretero-Guillén A, Márquez-Ruiz J: **Exploring new transcranial electrical stimulation strategies in animal models for brain function modulation**. *Curr Opin Biomed Eng* 2018.

Up to date review and outlook of research of tCS in animal models, with a discussion on emerging trends to improve focality and reach deeper targets.

49. Márquez-Ruiz J, Leal-Campanario R, Sánchez-Campusano R, Molaei-Ardekani B, Wendling F, Miranda PC, Ruffini G, Guart A, Delgado-García JM: **Transcranial direct-current stimulation modulates synaptic mechanisms involved in associative learning in behaving rabbits**. *PNAS* 2012, **109**.

50. Paciello F, Podda MV, Rolesi R, Cocco S, Petrosini L, Troiani D, Fetoni AR, Paludetti G, Grassi C: **Anodal transcranial direct current stimulation affects auditory cortex plasticity in normal-hearing and noise-exposed rats**. *Brain Stimul* 2018, **11**: 1008–1017. <https://doi.org/10.1016/j.brs.2018.05.017>. Epub 2018 May 31.

51. Wischnewski M, Engelhardt M, Salehinejad MA, Schutter DJLG, Kuo MF, Nitsche MA: **NMDA receptor-mediated motor cortex plasticity after 20 Hz transcranial alternating current stimulation**. *Cerebr Cortex* 2018. <https://doi.org/10.1093/cercor/bhy160>.

52. Fonteneau C, Redoute J, Haesebaert F, Le Bars D, Costes N, Suaud-Chagny MF, Brunelin J: **Frontal transcranial direct current stimulation induces dopamine release in the ventral striatum in human**. *Cerebr Cortex* 2018, **28**:2636–2646.

53. Santana-Gómez CE, Alcántara-González D, Luna-Munguía H, Bañuelos-Cabrera I, Magdaleno-Madrugal V, Fernández-Mas R, Besio W, Rocha L: **Transcranial focal electrical stimulation reduces the convulsive expression and amino acid release in the hippocampus during pilocarpine-induced status epilepticus in rats**. *Epilepsy Behav* 2015, **49**:33–39. <https://doi.org/10.1016/j.yebeh.2015.04.037>. Epub 2015 May 23.

54. Pugliatti M, Beghi E, Forsgren L, Ekman M, Sobocki P: **Estimating the cost of epilepsy in Europe: a review with economic modeling. Estimating the cost of epilepsy in Europe: a review with economic modeling**. *Epilepsia* 2007, **48**: 2224–2233.

55. Bartolomei F, Lagarde S, Wendling F, McGonigal A, Jirsa V, Guye M, Benar C: **Defining epileptogenic networks: contribution of SEEG and signal analysis**. *Epilepsia* 2017, **58**: 1131–1147. <https://doi.org/10.1111/epi.13791>.

This paper provides a historical overview of the emerging epileptogenic network concept, emphasizing the idea that “focal” epilepsies actually involve networks at varying scales, how to identify them, and the practical relevance of these findings to therapy.

56. Wendling F, Bartolomei F, Mina F, Huneau C, Benquet P: **Interictal spikes, fast ripples and seizures in partial epilepsies—combining multi-level computational models with**

**experimental data**. *Eur J Neurosci* 2012, **36**:2164–2177. <https://doi.org/10.1111/j.1460-9568.2012.08039.x>.

57. Hutchings F, Han CE, Keller SS, Weber B, Taylor PN, Kaiser M: **Predicting surgery targets in temporal lobe epilepsy through structural connectome based simulations**. *PLoS Comput Biol* 2015, **11**. e1004642–24.

58. Khambhati AN, Davis KA, Lucas TH, Litt B, Bassett DS: **Virtual cortical resection reveals push-pull network control preceding seizure evolution**. *Neuron* 2016, **91**:1170–1182.

59. Goodfellow M, Rummel C, Abela E, Richardson MP, Schindler K, Terry JR: **Estimation of brain network ictogenicity predicts outcome from epilepsy surgery**. *Sci Rep* 2016, **6**:1–13.

60. Proix T, Bartolomei F, Guye M, Jirsa VK: **Individual brain structure and modelling predict seizure propagation**. *Brain* 2017, **140**:641–654.

61. Sinha N, Dauwels J, Kaiser M, Cash SS, Brandon Westover M, Wang Y, Taylor PN: **Predicting neurosurgical outcomes in focal epilepsy patients using computational modelling**. *Brain* 2017, **140**:319–332.

In this work, authors predict surgical outcomes in epilepsy patients with high accuracy using personalized brain network models. They substitute the functional connectivity of the brain, derived from ECoG recordings of epileptic patients, for structural connectivity as a basis for the computational models.

62. Regner GG, Pereira P, Leffa DT, de Oliveira C, Vercelino R, Fregni F, Torres I: **Preclinical to clinical translation of studies of transcranial direct-current stimulation in the treatment of epilepsy: a systematic review**. *Front Neurosci* 2018, **12**:189. <https://doi.org/10.3389/fnins.2018.00189>.

A recent review of studies assessing the efficacy of tDCS in epilepsy.

63. Fox MD, Randy L, Buckner Hesheng Liu, Chakravarty MMallar, Lozano Andres M, Pascual-Leone Alvaro: **Linking invasive and noninvasive brain stimulation**. *Proc Natl Acad Sci USA* 2014b, **111**:E4367–E4375. <https://doi.org/10.1073/pnas.1405003111>.

64. Ota B, Dutta A, Foerster Á, Ripolles O, Kuceyeski A, Miranda PC, Edwards DJ, Ilić TV, Nitsche MA, Ruffini G: **Opportunities for guided multichannel non-invasive transcranial current stimulation in poststroke rehabilitation**. *Front Neurol* 2016, **7**:21. <https://doi.org/10.3389/fneur.2016.00021>.

This paper discusses potential opportunities for neuroimaging-guided tDCS-based rehabilitation strategies after stroke that could be personalized using brain networks.

65. Alstott J, Breakspear M, Hagmann P, Cammoun L, Sporns O: **Modeling the impact of lesions in the human brain**. *PLoS Comput Biol* 2009, **5**:e1000408–e1000412.

66. Aerts H, Fias W, Caeyenberghs K, Marinazzo D: **Brain networks under attack: robustness properties and the impact of lesions**. *Brain* 2016, **139**:3063–3083.

67. Corbetta M, Ramsey Lenny, Callejas Alicia, Baldassarre Antonello, Hacker Carl D, Siegel Joshua S, Astafiev Serguei V, Rengachary Jennifer, Zinn Kristina, Lang Catherine E, Connor Lisa Tabor, Fucetola Robert, Strube Michael, Carter Alex R, Shulman Gordon L: **Common behavioral clusters and subcortical anatomy in stroke**. *Neuron* 2015, **85**:927–941. <https://doi.org/10.1016/j.neuron.2015.02.027>. March 4.

68. Kuceyeski A, Navi BB, Kamel H, Raj A, Relkin N, Togliola J, Iadecola C, O'Dell M: **Structural connectome disruption at baseline predicts 6-months post-stroke outcome**. *Hum Brain Mapp* 2016, **37**:2587–2601. <https://doi.org/10.1002/hbm.23198>.

Structural connectome disruption is estimated using the *NeMo* tool developed by the authors (which allows estimation of connectome disruption from MRI). The measures at baseline predict 6-months post-stroke outcome in various functional domains including cognition, motor function, and daily activities.